

Prism™ SG

HERBICIDE

Outstanding post-emergent control of quackgrass, redroot pigweed and annual grasses.

WHY USE PRISM™ SG HERBICIDE?

- Can be used on all types of potatoes, including seed and early maturing varieties.
- Flexible re-cropping options.

BROADLEAF WEEDS CONTROLLED

- Redroot pigweed

GRASS WEEDS CONTROLLED

- Barnyard grass
- Fall panicum
- Green foxtail
- Witchgrass
- Yellow foxtail

PERENNIAL WEEDS CONTROLLED

- Quackgrass

BROADLEAF WEEDS SUPPRESSED

- Lamb's-quarters

CROP ROTATION

ANY TIME

- Field corn

FOUR MONTHS AFTER APPLICATION

- Winter wheat

10 MONTHS AFTER APPLICATION

- Barley
- Canola
- Chickpeas
- Corn (sweet or seed)
- Dry beans

- Durum wheat
- Faba beans
- Field peas
- Flax
- Lentils
- Oats
- Potatoes

- Red clover
- Sorghum
- Soybeans
- Spring wheat
- Sunflowers
- White beans

APPLICATION GUIDELINES

| | | |
|--------------------------------------|---------------------|---|
| <div>CROPS</div> <div>Potatoes</div> | PACKAGING | 12 x 480 g bottles |
| | RATES | 24 g/ac |
| | ACRES TREATED | 20 ac/bottle |
| | WATER VOLUME | 40 L/ac (10 US gal/ac) |
| | TIMING | <ul style="list-style-type: none">• Apply as a broadcast spray, with a recommended surfactant, to potatoes prior to initiation of flowering• Recommended non-ionic surfactant: Citowett Plus, Agral 90 or Ag-Surf at 2 L per 1000 L spray solution (0.2% v/v)• Application to control annual grasses and quackgrass must be made before the crop canopy can interfere with spray coverage of the target weeds. Cultivation is NOT recommended within 7-10 days prior to or after application. |
| | RAINFAST | 4 hours |
| | TANK-MIX HERBICIDES | Tricor 75DF |



DOWNLOAD THE 2025 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.

APPLICATION TIMING

Weeds which emerge after application of Prism™ SG herbicide will not be controlled. Application should be made when the majority of weeds have emerged. Annual grass and broadleaf weeds are most sensitive when small and actively growing. Early crop establishment and a good crop stand are important in providing competition for weeds and effective post-emergence control of quackgrass and annual weeds when using Prism™ SG.

PRE-HARVEST INTERVAL

30 days



Always read and follow label directions.