

Ares™ SN

The performance standard in weed control for Clearfield® canola.

HERBICIDE

WHY USE ARES™ SN HERBICIDE?

- Consistent and reliable post-emergent weed control, including subsequent flushes.
- Wide window of application on both crop and weeds.
- Superior control of lamb's quarters, wild buckwheat, cleavers and volunteer canola.

BROADLEAF WEEDS CONTROLLED

- Chickweed
- Cleavers^{2,3}
- Cow cockle
- Hemp-nettle²
- Lamb's-quarters²
- Redroot pigweed
- Round-leaved mallow

- Russian thistle
- Shepherd's purse²
- Smartweed²
- Stinkweed
- Stork's bill
- Wild buckwheat
- Wild mustard
- Volunteer tame mustard
- Volunteer canola (excluding Clearfield® varieties)

GRASS WEEDS CONTROLLED

- Barnyard grass
- Japanese brome¹
- Yellow foxtail
- Green foxtail
- Persian darnel
- Wild oats
- Volunteer canary seed
- Volunteer barley

- Volunteer tame oats
- Volunteer spring wheat (excluding Clearfield® varieties)
- Volunteer durum wheat

Surjet™

ADJUVANT

Surjet™ Adjuvant is a blend of surfactant and petroleum hydrocarbons designed for use with Ares™ SN or Amity WDG herbicides.

Rate: 0.5% v/v

Water volume: 40 L/ac (10 US gal/ac)

Surjet Adjuvant is sold in a co-pack with Ares SN.

¹ For spring germinating Japanese brome control apply at 1 to 4-leaf stage
² Group 2 susceptible biotypes will not be controlled

³ A tank mix with Facet® L will improve control of Group 2 resistant biotypes

APPLICATION GUIDELINES

CROPS Clearfield® Canola	PACKAGING	Case: • Ares™ SN: 1 x 9.8 L jug • Surjet Adjuvant: 1 x 8.1 L jug
	RATES & ACRES TREATED	40 ac/case
	WATER VOLUME	• Ground 40 L/ac (10 US gal/ac) • Aerial NOT registered
	TIMING	Crop stage: 2 to 7-leaf stage Weed stage: 1 to 6-leaf stage for grassy weeds and cotyledon to 4-leaf stage for broadleaf weeds
	RAINFAST	2 hours
	TANK MIX HERBICIDES	Lontrel™ XC

MIXING INSTRUCTIONS

1. Fill the spray tank with ½ to ¾ of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add required amount of Ares SN
4. Add any tank mix partners
5. Add the required amount of Surjet adjuvant at 0.5% v/v
6. Complete filling the sprayer tank with water

Note: Do not mix undiluted herbicides in the chem-handler

CROP ROTATION

Sensitivity of injury can vary depending on the crop. If rainfall in your area was less than 125 mm during the growing season between **June 1 to August 31**, it is recommended that you grow field peas, Clearfield® lentils or Clearfield® wheat on Clearfield® canola stubble.

The following crops may be grown safely the year following an application:

- Canary seed
- Chickpeas
- Field peas
- Field corn
- Clearfield® canola
- Clearfield® canola-quality Brassica juncea
- Lentils (including Clearfield® lentils)
- Spring wheat (including Clearfield® varieties)
- Spring barley

The following crops may be grown safely two years following an application:

- Canola
- Durum wheat
- Flax
- Sunflower

GRAZING AND HARVEST

Do not harvest treated crop within 60 days after application.