

# Korrex™ II

## HERBICIDE

Korrex™ II herbicide delivers superior pre-seed control of kochia, including Group 2 and 9 resistant biotypes, as well as 21 other tough broadleaf weeds.

### WHY USE KORREX™ II HERBICIDE?

- Kochia control. Superior pre-seed control of kochia, including Group 2 and Group 9 resistant biotypes.
- Flexible mixing options. Easy-mixing tank additive for any glyphosate.
- Broad-spectrum broadleaf weed control, including dandelion, narrow-leaved hawk's beard, wild buckwheat, flixweed and stinkweed.
- Extended control of volunteer canola flushes.
- Effective solution for herbicide resistance management.
- Excellent solution for Canada thistle control with a fall application.

### KORREX™ II + GLYPHOSATE AT 0.5 REL/AC

#### BROADLEAF WEEDS CONTROLLED

- Annual sow thistle<sup>2</sup>
- Canada fleabane<sup>3\*</sup>
- Chickweed\*
- Cleavers\*
- Common ragweed\*
- Cow cockle\*
- Flixweed\*
- Hemp-nettle\*
- Kochia\*
- Lamb's-quarters\*
- Narrow-leaved hawk's beard
- Redroot pigweed\*
- Russian thistle
- Scentless chamomile<sup>2</sup>
- Shepherd's purse\*
- Smartweed\*
- Stinkweed\*
- Volunteer canola<sup>\*1</sup>
- Volunteer flax
- Wild buckwheat\*
- Wild mustard\*

#### GRASS WEEDS CONTROLLED

- Downy brome
- Giant foxtail
- Green foxtail
- Persian darnel
- Volunteer barley
- Volunteer wheat
- Wild oats

#### PERENNIAL WEEDS CONTROLLED

- Dandelion\* (seedling, overwintered rosettes, mature plants up to 30 cm diameter)
- Perennial sow thistle<sup>4</sup>

#### SPRING AND FALL SOILACTIVE™ EXTENDED CONTROL

- Canada fleabane
- Canola (except Clearfield<sup>®</sup>)
- Chickweed<sup>5</sup>
- Cleavers<sup>5</sup>
- Common ragweed
- Dandelion seedling
- Flixweed
- Hemp-nettle<sup>5</sup>
- Lamb's-quarters<sup>5</sup>

- Narrow-leaved hawk's beard<sup>5</sup>
- Redroot pigweed
- Scentless chamomile
- Shepherd's purse<sup>5</sup>
- Smartweed<sup>5</sup>

- Stinkweed<sup>5</sup>
- Wild buckwheat
- Wild mustard

### KORREX™ II + GLYPHOSATE AT 1-2.8 REL/AC

- Annual sow thistle
- Canada thistle (rosette stage)
- Quackgrass

\*Weed controlled through multi-effective mode of action

1 Including all herbicide-tolerant canola varieties

2 Suppression only

3 Less than 8 cm in height

4 Applications made at advanced stages will reduce effectiveness

5 Will not provide extended control of Group 2 resistant biotypes

## APPLICATION GUIDELINES

<b>CROPS</b>  Prior to cereal crops (spring or fall application):  Barley  Durum wheat  Oats  Spring wheat  Winter wheat	<b>PACKAGING</b>	• Korrex II A: 1 x 0.45 kg jug • Korrex II B: 1 x 7.76 L jug
	<b>RATES</b>	<b>Spring Rate:</b> • Korrex II A: 5.7 g/ac • Korrex II B: 97 ml/ac  <b>Fall Rate:</b> • Korrex II A: 8.1 g/ac • Korrex II B: 139 ml/ac
	<b>ACRES TREATED</b>	<b>Spring:</b> 80 ac/case <b>Fall:</b> 56 ac/case
	<b>WATER VOLUME</b>	Ground 20-40 L/ac (5-10 gal/ac)
	<b>TIMING</b>	<b>Spring application:</b> Prior to seeding (no later than 48 hours after seeding) <b>Fall application:</b> From after harvest to freeze up
	<b>RAINFAST</b>	30 minutes
	<b>HERBICIDE TANK MIX HERBICIDES</b>	• VP480 • Compatible with all forms of glyphosate

### MIXING INSTRUCTIONS

1. Fill sprayer tank ½ full of water
2. Start sprayer tank agitation
3. Add the required amount of Korrex II A herbicide, continue agitation
4. Add the required amount of Korrex II B herbicide, continue agitation
5. Add the required amount of glyphosate, continue agitation
6. Fill the sprayer tank with sufficient water to spray 20-40 L/ac (5-10 US gal/ac)

Note: Do not mix undiluted herbicides in the chem-handler

### CROP ROTATION

The year following a spring Korrex II application, fields can be seeded to:

- Barley
- Canola
- Chickpeas
- Corn
- Field beans
- Flax
- Lentils
- Mustard
- Peas
- Potatoes (except seed potatoes)
- Soybeans
- Summerfallow
- Sunflower
- Wheat

### APPLICATION TIMING AND SEEDING

- Korrex II can be applied in the spring prior to seeding wheat, barley or oats and as an initial treatment in summerfallow.
- Korrex II applied in the fall (after August 1), can be seeded to winter wheat that fall or spring wheat, durum, barley and oats the following spring.
- Newly registered Spring Rate for use in the fall. Get more acres of Korrex™ II without compromising control, spray early and save.

### GRAZING AND HARVEST

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest treated crop within 60 days after application.