

Proactive resistance management with rotational flexibility.

WHY USE OnDeck™ HERBICIDE?

- Provides control of key weeds like kochia, wild mustard, wild buckwheat, and volunteer canola
- Additional control of green and yellow foxtail without a Group 1 or 2 herbicide
- Two different modes of action built to provide proactive resistance management
- Complete rotational freedom

ONDECK™ ALONE WEEDS CONTROLLED

BROADLEAF

- Chickweed
- Cleavers (up to 4 whorl)
- Hemp-nettle
- Kochia²
- Smartweed
- Lamb's quarters
- Redroot pigweed
- Volunteer canola
- Wild buckwheat (up to 4 leaf)
- Wild mustard

GRASS

- Green foxtail¹
- Yellow foxtail

When OnDeck™ is tank mixed with Simplicity™ GoDRI™ (full rate) these additional weeds are controlled or suppressed:

GRASS

- Wild oats
- Japanese brome
- Barnyard grass

BROADLEAF

- Corn spurry
- Cow cockle
- Flixweed
- Round-leaved mallow
- Stinkweed

WEEDS SUPPRESSED

- Persian darnel
- Downy brome
- Canada thistle
- Dandelion
- Russian thistle
- White cockle

¹ Including Group 1 and Group 2 resistant biotypes

² Group 2, Group 4, and Group 9 resistant biotypes

APPLICATION GUIDELINES

CROP

Spring wheat
Durum wheat
Winter wheat
Barley

PACKAGING

Case: 2 x 6.475L jugs
Tote: 414.4L

RATE

Standard Rate (Recommended): 324 ml/ac
High Rate: 435 ml/ac

ACRES TREATED

Standard Rate (Recommended)
20ac/jug, 40 ac/case, 1280 ac/tote

High Rate
15ac/jug, 30 ac/case, 956 ac/tote

CROP

Ground: 10 US gal/ac
Range: 5-20 US gal/ac
Aerial: minimum 3 US gal/ac

Add 0.5% v/v adjuvant (MSO, COC or HSOC) for improved control of larger weeds or when weeds are under environmental stress

TIMING

Crop stage: 1st leaf to jointing (1st node visible, BBCH31)
Weed stage: 1 to 4 leaf
Rainfast: 1 hour

TANK MIX HERBICIDES

- Simplicity™ GoDRI™
- Simplicity™
- Trondus® Herbicide*
- Axial® Herbicide*
- Horizon® NG Herbicide
- Everest® 2.0 Herbicide**
- MCPA Ester 600
- 2,4-D Ester 700

*Axial or Trondus Herbicide may be substituted with any equivalent pinoxaden product

**Everest 2.0 may be substituted with any equivalent flucarbazone product

CROP ROTATION

IMMEDIATE RE-PLANT OPTIONS

Corn (sweet corn, seed corn, and popcorn)
Barley
Wheat (spring, durum, and winter)

10 MONTHS AFTER APPLICATION

Alfalfa	Lentils
Canola	Oats
Dry bean	Mustard (oriental, brown, and yellow)
Fall Rye	Potato (not for seed)
Peas (field and edible)	Soybean
Grass grown for seed or forage	Sunflower

GRAZING AND HARVEST

Allow 30 days after application before grazing, cutting hay, or silage.

Mature crops may be harvested 50 days after application.