

Pixxaro™

Arylex™ active

HERBICIDE

Puts you in control and provides the flexibility to tank mix with ANY graminicide partner.

WHY USE PIXXARO™ HERBICIDE?

- Delivers exceptional broadleaf weed control in wheat and barley.
- Contains Arylex™ active which allows farmers to Just GO on small or large weeds, early or late crop staging, and even in cool or dry conditions.
- Provides flexibility to tank mix with the right graminicide partner.

BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Annual sow thistle
- Annual sunflower
- Ball mustard
- Burdock
- Canada fleabane
- Chickweed¹
- Cleavers¹
- Cocklebur
- Common ragweed
- Flixweed

- Hemp-nettle¹
- Henbit
- Kochia²
- Lamb's-quarters
- Night shade species (*Eastern black, hairy and cutleaf*)
- Plantain
- Prickly lettuce
- Ragweed (*false and giant*)
- Redroot pigweed
- Round-leaved mallow
- Shepherd's purse

- Stinkweed
- Stork's-bill
- Velvetleaf
- Vetch
- Volunteer alfalfa
- Volunteer canola (*all herbicide tolerant biotypes*)
- Volunteer flax
- Wild buckwheat
- Wild mustard¹
- Wild radish

GRASS WEEDS CONTROLLED

- Barnyard grass

WEEDS SUPPRESSED

- Canada thistle
- Dandelion
- Field horsetail
- Perennial sow thistle
- Smartweed

For a complete weed list and specific weed staging, please refer to the Pixxaro label.

¹ Including Group 2 resistant biotypes
² Including Group 2 and Group 9 resistant biotypes

APPLICATION GUIDELINES

CROPS Barley Durum wheat Spring wheat Winter wheat	PACKAGING Case: • Pixxaro A: 1 x 4.9 L jug • Plus M Ester 600: 1 x 9.45 L jug Bulk Pack: • Pixxaro A: 4 x 9.8 L jugs • Plus M Ester 600: 1 x 75.1 L drum Pallet: • Pixxaro A: 16 x 9.8 L jugs • Plus M Ester 600: 4 x 75.1 L drums
	RATES • Pixxaro A: 125 ml/ac • Plus M Ester 600: 235 ml/ac
	ACRES TREATED • 40 ac/case • 320 ac/bulk pack
	WATER VOLUME Ground 20-40 L/ac (5-10 US gal/ac)
	TIMING Crop stage: 3-leaf to just prior to flag leaf emergence Weed stage: 1 to 8-leaf (or larger; see label)
	RAINFAST 1 hour
	TANK MIX HERBICIDES • Axial® • Everest® • Horizon • Liquid Achieve™ • Puma® Advance • Simplicity™ • Simplicity™ GoDRI™ • Traxos



DOWNLOAD THE 2023 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

- Spring wheat
- Barley
- Oats
- Canola
- Corn
- Soybeans
- Sunflowers
- Faba beans
- Flax
- Field peas
- Potatoes (*except seed potatoes*)
- Mustard
- Alfalfa
- Dry bean (*Phaseolus vulgaris species including pinto, kidney and white types*)
- Timothy

GRAZING AND PRE-HARVEST INTERVAL

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.
- Do not cut the treated crop for hay or silage within 21 days after application.

AVAILABLE IN BULK