CEREAL PRE-SEED FLORASULAM: 2 g/ac 2

PrePass FLEX

Transform your glyphosate.

GROUP

HERBICIDE

WHY USE PREPASS™ FLEX HERBICIDE?

- · Flexibility to mix with any glyphosate formulation, at the rate of glyphosate you choose.
- · Advanced dry formulation that disperses quickly and completely for easy mixing
- · Convenient packaging 640 acre case. No heated storage required.
- · SoilActive™ technology for extended broadleaf weed control.
- Time management. Get the work done earlier spring or fall.

PREPASS™ FLEX HERBICIDE + GLYPHOSATE AT 0.5 REL/AC

BROADLEAF **WEEDS** CONTROLLED

- · Annual sow thistle1
- · Canada fleabane
- Chickweed
- Cleavers
- Common ragweed
- Cow cockle Flixweed
- · Hemp-nettle
- Kochia
- · Lamb's-quarters
- · Narrow-leaved hawk's beard

· Redroot pigweed · Russian thistle

- Scentless
- chamomile
- · Shepherd's purse
- Smartweed
- Stinkweed
- Volunteer canola³
- · Volunteer flax
- Wild buckwheat
- · Wild mustard

GRASS WEEDS CONTROLLED

- · Downy brome
- · Foxtail barley
- Giant foxtail
- · Green foxtail
- · Persian darnel
- Volunteer barley
- · Volunteer wheat
- · Wild oats

· Wild buckwheat · Wild mustard

PERENNIAL WEEDS

CONTROLLED

· Dandelion²

sow thistle1

· Quackgrass¹

Perennial

· Canada thistle

SPRING AND FALL SOILACTIVE™ **EXTENDED** CONTROL:

- · Canada fleabane
- · Chickweed4
- · Cleavers4
- · Common ragweed
- · Dandelion seedling · Redroot pigweed
- Flixweed
- · Hemp-nettle⁴ · Lady's-thumb
- · Lamb's-quarters4
- · Narrow-leaved
- hawk's beard⁴

- · Smartweed⁴

chamomile

Scentless

- · Stinkweed⁴
- · Volunteer canola (except Clearfield®)

· Shepherd's purse

1 Requires elevated rate of glyphosate; refer to glyphosate label

APPLICATION GUIDELINES

	PACKAGING	8 x 0.648 kg jugs
CROPS Prior to cereal crops (spring or fall application): Barley Durum wheat Oats Spring wheat Winter wheat	RATES	Spring & Fall Rate: 8.1 g/ac
	ACRES TREATED	80 ac/jug (640 ac/case)
	WATER VOLUME	Ground 20-40 L/ac (5-10 US gal/ac)
	TIMING	Spring application: Prior to seeding (no later than 48 hours after seeding) Chemfallow: When weeds are actively growing, 1 to 4-leaf stage Fall application: From after harvest to freeze up
	RAINFAST	30 minutes
	TANK MIX HERBICIDES	VP480 Compatible with all forms of glyphosate

MIXING INSTRUCTIONS

- 1. Fill sprayer tank ½ full of water
- 2. Start sprayer tank agitation
- 3. Add the required amount of PrePass FLEX herbicide, continue agitation
- 4. Add the required amount of glyphosate, continue agitation
- 5. Fill the sprayer tank with sufficient water to spray 20-40 L/ac (5-10 US gal/ac)

CROP ROTATION

The year following a spring PrePass FLEX application, fields can be seeded to:

- Alfalfa · Faba beans Oats Summerfallow Barlev · Field beans · Peas Sunflower · Canola Flax Potatoes Wheat (except seed potatoes) Chickpeas Lentils
- Soybeans • Corn Mustard

APPLICATION TIMING AND SEEDING

- PrePass FLEX can be applied in the spring prior to planting wheat, barley or oats.
- PrePass FLEX applied after August 1 can be seeded to winter wheat that fall or spring wheat, durum, barley and oats the following spring.

GRAZING AND HARVEST

- · Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.

36 37

² Seedling, overwintered rosettes, mature plants up to 30 cm in diameter

³ All herbicide-tolerant varieties

⁴ Will not provide extended control of Group 2 resistant biotypes