

# Prestige™ XL

Maximize potential.  
Every acre, clean.

## HERBICIDE

### WHY USE PRESTIGE™ XL HERBICIDE?

- Excellent control of thistles, cleavers, kochia and more.
- Your cleanest cereal crops, year after year.
- Multiple pack sizes: cases and drums.

#### 950 ml/ac (HIGH RATE)

<b>BROADLEAF WEEDS CONTROLLED</b>	• Field horsetail <sup>5</sup> • Flixweed • Hemp-nettle • Annual sow thistle • Annual sunflower • Burdock • Canada thistle <sup>1</sup> • Chickweed <sup>3, 4</sup> • Cleavers <sup>3</sup> • Cocklebur • Common groundsel	• Kochia <sup>3</sup> • Lamb's-quarters • Plantain <sup>5</sup> • Prickly lettuce • Ragweed • Redroot pigweed • Round-leaved mallow	• Russian pigweed • Scentless chamomile • Shepherd's purse • Smartweed • Stinkweed • Stork's-bill • Tartary buckwheat • Vetch • Volunteer canola <sup>6</sup> • Volunteer flax	• Volunteer sunflower • Wild buckwheat • Wild mustard • Wild radish
-----------------------------------	--	---	---	--

#### PERENNIAL WEEDS CONTROLLED

- Canada thistle<sup>1</sup>
- Dandelion<sup>2</sup>
- Perennial sow thistle<sup>1</sup>

#### 710 ml/ac (LOW RATE)

<b>BROADLEAF WEEDS CONTROLLED</b>	• Flixweed • Kochia <sup>3</sup> • Lamb's-quarters • Plantain <sup>5</sup> • Prickly lettuce • Ragweed • Shepherd's purse • Stinkweed	• Stork's-bill • Vetch • Volunteer flax • Volunteer sunflower • Wild buckwheat • Wild mustard • Wild radish	<b>PERENNIAL WEEDS CONTROLLED</b>	• Canada thistle <sup>1</sup> (low infestations)
-----------------------------------	--	---	-----------------------------------	---

#### WEEDS SUPPRESSED

- Volunteer canola<sup>6</sup>

## APPLICATION GUIDELINES

<b>CROPS</b> Barley Canary seed Durum wheat Forage grasses (see label for complete list.) Oats Spring wheat Winter wheat	<b>PACKAGING</b>	Case: 2 x 9.5 L jugs Drum: 113.6 L drum
	<b>RATES</b>	Low rate: 710 ml/ac High rate: 950 ml/ac
	<b>ACRES TREATED</b>	Low rate: • 13.5 ac/jug • 160 ac drum High rate: • 10 ac/jug • 120 ac/drum
	<b>WATER VOLUME</b>	• Ground 20-40 L/ac (5-10 US gal/ac) • Aerial 12-20 L/ac (3-5 US gal/ac)
	<b>TIMING</b>	Crop stage: 3-leaf to just prior to flag leaf emergence Weed stage: 1 to 6-leaf stage
	<b>RAINFAST</b>	4 hours
	<b>TANK MIX HERBICIDES</b>	• Axial® • Assert® • Everest® • Horizon • Liquid Achieve™ SC • Puma® • Simplicity™ • Simplicity™ GoDRI™



DOWNLOAD THE 2023 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

### CROP ROTATION

- Barley
- Canola
- Flax
- Mustard
- Oats
- Peas<sup>7</sup>
- Rye
- Sugar beets
- Summerfallow
- Wheat

### GRAZING AND HARVEST

- Livestock may graze treated areas 7 days after application.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 7 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.

<sup>7</sup> For pea or soybean rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140 mm (5.5 inches) and annual rainfall must be greater than 175 mm (6.9 inches).

<sup>1</sup> Season-long control, with some regrowth in the fall (top growth control)

<sup>2</sup> Spring rosettes only

<sup>3</sup> Including Group 2 resistant biotypes

<sup>4</sup> Controls or suppresses the population that is present at time of spraying but not subsequent flushes

<sup>5</sup> Top growth only

<sup>6</sup> All herbicide-tolerant canola varieties