

Prism™ SG

Outstanding post-emergent control of quackgrass, pigweed and annual grasses.

HERBICIDE

WHY USE PRISM™ SG HERBICIDE?

- Can be used on all types of potatoes, including seed and early maturing varieties.
- Flexible re-cropping options.

BROADLEAF WEEDS CONTROLLED

- Hairy nightshade
- Redroot pigweed

GRASS WEEDS CONTROLLED

- Barnyard grass
- Fall panicum
- Green foxtail
- Witchgrass
- Yellow foxtail

CROP ROTATION

10 MONTHS AFTER APPLICATION

- Barley
- Canola
- Chickpeas
- Corn (*sweet or seed*)
- Dry beans
- Faba beans
- Field peas
- Flax

PERENNIAL WEEDS CONTROLLED

- Quackgrass

WEEDS SUPPRESSED

- Lamb's-quarters

FOUR MONTHS AFTER APPLICATION

- Winter wheat

ANY TIME

- Field corn

- Lentils
- Oats
- Potatoes
- Red clover
- Sorghum
- Soybeans
- Spring wheat (*including durum*)
- Sunflowers
- White beans

APPLICATION GUIDELINES

CROPS Potatoes	PACKAGING	12 x 480 g bottles
	RATES	24 g/ac
	ACRES TREATED	20 ac/bottle
	WATER VOLUME	40 L/ac (10 US gal/ac)
	TIMING	<ul style="list-style-type: none"> • Apply as a broadcast spray, with a recommended surfactant, to potatoes prior to initiation of flowering • Recommended non-ionic surfactant: Citowett Plus, Agral 90 or Ag-Surf at 2 L per 1000 L spray solution (0.2% v/v) • Application to control annual grasses and quackgrass must be made before the crop canopy can interfere with spray coverage of the target weeds. Cultivation is NOT recommended within 7-10 days prior to or after application.
	RAINFAST	4 hours
TANK MIX HERBICIDES	Tricor 75DF	

MIXING INSTRUCTIONS

1. Fill clean tank about 1/3 full of clean water
2. With the agitator running, add the required amount of Prism™ SG herbicide. Continue to agitate for a minimum of 5 minutes to ensure that Prism SG is completely dissolved
3. Make sure there are no un-dissolved granules, and then add the required amount of the tank mix partner
4. After Prism SG has been well mixed and is in suspension, add a recommended non-ionic surfactant such as Citowett Plus, Agral 90, or Ag-Surf at 2 L per 1000 L spray solution (0.2% v/v)
5. Fill the remainder of the spray tank with water

APPLICATION TIMING

Weeds which emerge after application of Prism SG will not be controlled. Application should be made when the majority of weeds have emerged. Annual grass and broadleaf weeds are most sensitive when small and actively growing. Early crop establishment and a good crop stand are important in providing competition for weeds and effective post-emergence control of quackgrass and annual weeds when using Prism SG.

PRE-HARVEST INTERVAL

30 days

Always read and follow label directions.