

Prominex™

Arylex™ active

HERBICIDE

Prominex™ delivers exceptional control of annual AND perennial broadleaf weeds with the convenience of an all-in-one formulation.

WHY USE PROMINEX HERBICIDE IN WHEAT AND BARLEY?

- **Unmatched Broadleaf Weed Control.** Prominex provides control of both annual AND perennial broadleaf weeds including Canada fleabane and Canada thistle.
- **Just GO.** Thanks to Arylex active, you can Just GO on small or large broadleaf weeds, in early or late crop staging, and even in cool or dry conditions.
- **Tank Mix Flexibility.** Tank-mix with MCPA Ester 600 or 2,4-D Ester for expanded Group 4 broadleaf weed control.
- **Convenient Formulation.** All-in-one ME formulation allows for ultimate convenience and easy loading.

Prominex + 235 ml/ac (5 oz./ac) MCPA Ester 600

GRASS WEEDS CONTROLLED

- Barnyard grass

ANNUAL BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Annual sow thistle
- Ball mustard
- Burdock
- Canada fleabane**
- Canada thistle
- Chickweed**
- Cleavers**
- Cocklebur
- Common ragweed**
- Cow cockle
- Dandelion

- False ragweed
- Flixweed*
- Giant ragweed**
- Hemp-nettle*
- Henbit
- Kochia**
- Lamb's-quarters
- Nightshade species (Eastern black, hairy and cutleaf)
- Plantain
- Prickly lettuce
- Redroot pigweed
- Round-leaved mallow
- Shepherd's purse*
- Stinkweed

- Stork's-bill
- Velvetleaf
- Vetch
- Volunteer alfalfa
- Volunteer canola (all herbicide tolerant varieties)
- Volunteer flax
- Wild buckwheat
- Wild mustard
- Wild radish

WEEDS SUPRESSED

- Canada thistle
- Field horsetail
- Perennial sow-thistle
- Smartweed

*Including Group 2 resistant biotypes

**Including Groups 2 & 9 resistant biotypes

For a complete weed list and specific weed staging, please refer to the Prominex RA label.

PROMINEX APPLICATION GUIDELINES

CROPS

Barley
Durum wheat
Spring wheat
Winter wheat

RATES AND ACRES TREATED

RATES:

– 414 ml/ac

ACRES TREATED:

– 40 ac/case (20 ac/jug)

PACKAGING:

– Case: 2 x 8.3 L jugs

WATER VOLUME:

– 10–20 US gal/ac

WHEN TO APPLY

• 3-leaf to just prior to flag leaf emergence

RAINFAST

• 4 hours

HERBICIDE TANK MIXES

• MCPA Ester 600
• 2, 4-D Ester 700
• Simplicity™ GoDRI

• Axial®
• Axial + MCPA Ester 600
• Liquid Achieve™ SC
• Puma® Advance

Application information

For the most complete spectrum of broadleaf weeds controlled, tank-mix Prominex with 235 ml/ac of MCPA Ester 600

Crop rotation

- Barley
- Canola
- Corn
- Flax
- Forage grasses
- Oats
- Mustard
- Peas¹
- Fall rye (*not underseeded with legumes, clover or alfalfa*)
- Soybeans¹
- Summerfallow
- Wheat

Pre-harvest interval

- The PHI for treated crops is 60 days
- The PHI for hay or silage is 21 days
- Livestock may be grazed on treated crops 7 days following application

¹ For pea or soybean rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140 mm (5.5 inches) and annual rainfall must be greater than 175 mm (6.9 inches).

2020 Research Authorizations

Prominex was showcased in 53 x 80 acre research authorization sites applied by farmers throughout Alberta, Saskatchewan and Manitoba.



Prominex + Simplicity GODRI
(10 Days after application)