

Simplicity™ GoDRI™

HERBICIDE

Superior performance, including elite Group 2 wild oat and bonus broadleaf weed control with no re-cropping restrictions in a convenient, easy-to-use GoDRI formulation.



WHY USE SIMPLICITY™ GODRI™ HERBICIDE?

- Superior performance. Elite grass and broadleaf weed control.
- Wide window of application. The only Group 2 wild oat product that can be applied up to flag leaf.
- Resistance management. Controls Group 1 resistant wild oats.
- Broadleaf herbicide tank mix partner flexibility.
- Rotational freedom to all major crops the following year.
- Convenient GoDRI™ herbicide formulation for fast, easy mixing and handling.

BROADLEAF WEEDS CONTROLLED

- Chickweed
- Cleavers*
- Corn spurry
- Cow cockle
- Flixweed
- Hemp-nettle
- Redroot pigweed
- Round-leaved mallow
- Shepherd's purse
- Smartweed*
- Stinkweed
- Volunteer canola (excluding Clearfield®)

GRASS WEEDS CONTROLLED

- Wild oats
- Barnyard grass
- Downy brome (fall application)
- Japanese brome
- Yellow foxtail

WEEDS SUPPRESSED

- Canada thistle
- Dandelion
- Downy brome (spring application)
- Green foxtail¹
- Persian darnel
- Russian thistle
- White cockle
- Wild buckwheat

WHEN TANK MIXING SIMPLICITY™ GODRI™ WITH A BROADLEAF HERBICIDE THAT DOES NOT REQUIRE A SURFACTANT, BINDEM UTILITY MODIFIER MUST BE USED. BINDEM IS SOLD SEPARATELY.

When spraying Simplicity™ GoDRI™ alone, the addition of a non-ionic surfactant is ALWAYS required. The following non-ionic surfactants can be used: Agral 90 at 0.25% v/v, Sentry at 0.25% v/v, Ag-Surf® Original at 0.25% v/v

*Not including group 2 resistant
¹ Corteva Agriscience research trials indicate that application to small stage, actively growing plants provides an increased level of control

APPLICATION GUIDELINES

| | | | |
|---|--|---|--|
| CROPS Durum wheat Spring wheat Winter wheat | PACKAGING | 4 x 2.24 kg jugs | |
| | RATES | 28 g/ac | |
| | ACRES TREATED | 80 ac/jug (320 ac/case) | |
| | WATER VOLUME | • Ground 12-40 L/ac (3-10 US gal/ac) • Aerial minimum 12 L/ac (3 US gal/ac) | |
| | TIMING | Crop stage: 2-leaf to flag leaf Weed stage: • 1 to 4-leaf, two tillers stage of wild oats • 1 to 5-leaf stage of broadleaf weeds | |
| | RAINFAST | 2 hours | |
| | TANK MIX HERBICIDES³ | • Attain™ XC • Buctril® M • Cirpreme™ XC • Curtail® M • Exhilarate™ • MCPA Ester 600 • OcTTain™ XL | • Prestige™ XL • Prominex™ • Pixxaro™ • Refine Extra® • Stellar™ XL • Thumper® • Tilt® fungicide |



DOWNLOAD THE 2023 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

- Alfalfa
- Barley
- Brown mustard
- Canola
- Canola-quality brassica juncea
- Chickpea
- Dry bean (*Phaseolus vulgaris* species)
- Field corn
- Field peas
- Flax
- Lentils
- Oats
- Potatoes
- Soybean
- Spring wheat
- Sunflower
- Yellow mustard

GRAZING AND HARVEST

- Livestock may graze treated areas 7 days after application.
- Mature crops may be harvested 50 days after application.

³ Unless otherwise noted. Always read and follow label directions.