# **BERRIES**

### PEST TIMING CALENDAR



Corteva Agriscience™ offers an outstanding lineup of customer-focused products for berry growers that fit well into an Integrated Pest Management (IPM) program.

### Success<sup>™</sup>

Qalcova active

#### **INSECTICIDE**

Success™ insecticide with Qalcova™ active (spinosad) is a proven insecticide of natural origin that helps producers grow high-quality berry crops. Success provides trusted control of a broad range of pests including blueberry flea beetle, oblique-banded leafroller and spotted wing drosophila with minimal impact to the environment and beneficial insects when used according to the label.

### **Delegate**<sup>™</sup>

Jemvelva" active

#### **INSECTICIDE**

Delegate™ insecticide with Jemvelva™ active (spinetoram) is a naturally-derived product that is fast-acting against spotted wing drosophila, oblique-banded leafroller and blueberry spanworm with minimal impact on beneficials and the environment. Its unique profile makes it an excellent tool for IPM programs for berry growers.

### **Entrust**<sup>™</sup>

Qalcova"active

#### **INSECTICIDE**

Entrust™ insecticide with Qalcova™ active (spinosad) is a proven natural insecticide for organic berry growers. Entrust has excellent activity on thrips, leafrollers, blueberry flea beetle and spotted wing drosophila. Entrust is a perfect fit for IPM programs.

## Intrepid™

### **INSECTICIDE**


Intrepid™ insecticide is a novel group 18 insecticide that controls pests at early stages to get your berries off to a strong start. It induces a premature molt to control lepidopterous pests which stops feeding within 24 hours. Intrepid delivers both ovicidal and larvicidal activity providing residual control for up to 10 to 14 days after application.

#### PRE-HARVEST INTERVALS IN BERRIES

(Refer to each product label for crop specific pre-harvest intervals)

Closer™ insecticide with Isoclast™ active	1 day	Delegate™ insecticide	1-3 days	Entrust™ insecticide	1-3 days
Success™ insecticide	1-3 days	Fontelis™ fungicide	0 days	Indar™ fungicide	30 days
Nova™ fungicide	1-14 days	Tanos™ fungicide	9 days	Lontrel™ XC herbicide	45 days to 10 months
		Prism™ SG herbicide	21 days to 14 months		





<sup>\*</sup> Suppression

This chart is intended for reference only and not intended to provide information for application. Always read and follow label directions. Chart shows possible pest timing, not spray timing.

For more information visit us at **Horticulture.corteva.ca** to find your Corteva Agriscience™ Horticulture Expert.



<sup>&</sup>lt;sup>1</sup> Apply only when bees are not visiting the area.