



with RINSKOR™ ACTIVE

Milestone™ NXT Herbicide

GROUP	4	HERBICIDE
-------	---	-----------

Selective herbicide for post-emergent control of annual and perennial broadleaf weeds, invasive plants and shrubs on rangeland, permanent pasture, rights-of way, industrial and other non-crop areas of Canada.

COMMERCIAL

WETTABLE GRANULES

READ THE LABEL AND THIS BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Aminopyralid	60.00 %
(present as potassium salt)	
Florpyrauxifen	4.77%
(present as benzyl ester)	

REGISTRATION NO. 34728 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.1 kg - bulk

Corteva Agriscience Canada Company

Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9
1-800-667-3852

™ Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

PERSONAL PROTECTIVE EQUIPMENT

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab and/or cockpit.

OPERATOR USE PRECAUTIONS

- Wash hands before eating, drinking, using tobacco or using the washroom.
- If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing.
- Remove personal protective equipment immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.

AERIAL APPLICATION PRECAUTIONS

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. The pilot is allowed to load premixed chemicals with a closed system.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

RESTRICTED ENTRY INTERVALS

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application on agricultural areas.

For rights-of-way, industrial and other non-crop areas, DO NOT enter or allow worker entry into treated areas until sprays have dried.

SPRAY DRIFT PRECAUTIONS

Apply only when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE

Store this product away from food or feed.

DISPOSAL

Recyclable containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Returnable containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

Milestone™ NXT Herbicide, applied post emergent, controls broadleaf weeds, invasive plants and woody plants in rangeland, permanent pasture, industrial areas (including but not limited to rights-of-way and military bases) and other non-crop areas. Applications should be avoided under conditions of drought or other environmental stresses. For more information, contact your local Corteva Agriscience representative.

For best results, apply Milestone NXT Herbicide to young weeds that are actively growing at time of application.

Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or www.corteva.ca for information before applying any tank mix that is not specifically recommended on this label.

DIRECTIONS FOR USE

READ ALL DIRECTIONS CAREFULLY BEFORE APPLYING. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

GENERAL USE PRECAUTIONS

Milestone NXT Herbicide is selective and highly active against annual and perennial broadleaf weeds, invasive plants and shrubs. This product is recommended for use on areas where loss of broadleaf forage plants, including legumes, can be tolerated. Injury to or loss of desirable plants may result unless the following precautions are observed. Do not apply or drain or flush equipment on or near desirable vegetation or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots. Do not apply on walks, driveways, tennis courts or other similar areas.

- Do not exceed 200 g/ha per year. Only one application per year is permitted.
- Milestone NXT Herbicide cannot be applied on domestic or commercial turf grass.

Sensitive Plants

Milestone NXT Herbicide works primarily through uptake by plant foliage and translocation throughout the plant. However, secondary herbicide activity may occur through soil uptake of the aminopyralid component of Milestone NXT Herbicide. Very small amounts of Milestone NXT Herbicide can kill or damage sensitive broadleaf plants. Only spray if injury to existing forage can be tolerated. Care should be taken to avoid spraying desirable broadleaved plants, during both growing and dormant periods.

Avoid application within the drip line (outermost edge of the tree canopy) of desirable coniferous and deciduous trees unless injury can be tolerated. Use special caution when using aminopyralid-based products around species with extensive lateral root systems, shallow rooting species, and those that propagate vegetatively through layering.

Clippings or hay from vegetation which has been treated with aminopyralid should not be used for composting or mulching. Aminopyralid residues pass through animals unchanged and are still herbicidally active. Do not transfer livestock from treated grazing areas onto broadleaf crop areas without first allowing 3 days of grazing on untreated grass pasture.

Hay cut from treated grass, in the year of application, should only be used on-farm to feed livestock. The manure or compost from animals fed the treated hay or grazing treated areas should only be used on registered use sites where the loss of broadleaf forage plants, including legumes, can be tolerated. Contact Corteva Agriscience for additional information on sensitive broadleaf plant species.

INDUSTRIAL AREAS (INCLUDING RIGHTS-OF-WAY), OTHER NON-CROP AREAS, RANGELAND AND PERMANENT PASTURE

APPLICATION METHODS

Milestone NXT Herbicide can be applied post-emergence as a broadcast spray (ground or aerial) or as a spot/strip application to control weeds. Post emergence applications should be made before bud stage or early flowering, unless otherwise specified. Best results are obtained when the spray volume is sufficient to provide uniform coverage of treated plants. For optimum uptake and translocation of the herbicide, avoid mowing, haying, shredding or disturbing treated areas for at least 7 days following application.

MILESTONE NXT HERBICIDE ALONE

Apply Milestone NXT Herbicide at a rate of 100 – 200 g/ha. Apply when targeted undesirable vegetation is actively growing. Only weeds and brush present at the time of application will be controlled. Apply with a minimum of 100 L spray volume/ha, which will assure uniform coverage. For better coverage use 200 litres/ha for ground application. Use a surfactant, such as Gateway Adjuvant or an NIS at 0.25-0.5% v/v or methylated seed oil (MSO) at 0.5-1% v/v.

Individual Plant or Spot/Strip Applications

Use the following mixing chart to mix Milestone NXT with enough water and surfactant to make the correct amount of spray solution. Use equipment designed for individual plant or spot/strip applications, such as a backpack sprayer or hose and handgun, to spray susceptible invasive plants Thoroughly and uniformly wet the foliage of all target plants but not to the point of runoff. Direct spray away from aquatic habitats and non-target terrestrial plants. DO NOT apply this product directly to fresh water habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), estuaries or marine habitats.

Mixing Chart

Milestone NXT Herbicide	Surfactant (at 0.5% v/v)*	Spray Solution (water and herbicide)
1.0-2.0 grams	50 mL	10 litres
10- 20 grams	500 mL	100 litres
100- 200 grams	5000 mL	1000 litres

*Surfactant such as Gateway Adjuvant, MSO, NIS (such as Ag-Surf, Agral 90 or Citowett Plus)

Use the higher rate in the rate range when growing conditions are less than favourable or when the plant foliage is tall and dense.

PREHARVEST/GRAZING INTERVALS

The following restrictions for livestock or lactating dairy animals grazing in treated areas must be followed:

- Forage: Grazing/feeding is permitted on the day of application (0 days)
- Hay: Allow 7 days after application to cut or feed hay.
- Allow 3 days of grazing on an untreated pasture (or feed untreated hay) before transferring livestock to areas where sensitive broadleaf crops may be grown.

Weeds Controlled or Suppressed by Milestone NXT Herbicide Alone

Rate of Milestone NXT Herbicide	Weeds controlled	Weeds suppressed
100 g/ha	Canada fleabane ¹ velvetleaf ² common waterhemp ³	Canada thistle spotted knapweed
117 g/ha	common ragweed yellow star thistle ⁴ plumeless thistle musk or nodding thistle Canada fleabane horsenettle perennial sowthistle oxeye daisy ⁶	western ragweed curly dock Canada goldenrod scentless chamomile sulphur cinquefoil ⁵

	tall buttercup Canada thistle spotted knapweed common broomweed bull thistle tropical soda apple ⁷ hairy buttercup tropic croton	
150 g/ha	Weeds listed above plus: cudweed curly dock western ragweed scentless chamomile bitter sneezeweed hairy fleabane tansy ragwort clover poison hemlock	absinth (wormwood) Canada goldenrod
200 g/ha	Weeds listed above plus: absinth (wormwood) prickly lettuce Fuller's teasel tall ironweed fireweed purple loosestrife orange hawkweed ¹¹ mullein rush skeletonweed Scotch thistle wild caraway cleavers ¹² common lamb's quarter ¹³ annual sowthistle ¹⁴ wild chervil ¹⁵ wild parsnip	Canada goldenrod common yarrow common tansy diffuse knapweed ⁸ dandelion Russian knapweed ⁹ Japanese knotweed ¹⁰

¹Canada fleabane: Bolting up to 20 cm

²Velvetleaf: Up to 3 leaf and 8 cm

³Common waterhemp: Up to 3 leaf and 8 cm

⁴Yellow star thistle: Apply to plants at the rosette through bolting growth stage

⁵Sulfur cinquefoil: apply to plants in the prebud stage of development.

⁶Oxeye daisy: Apply to plants in the pre-bud stages of development

⁷Tropical soda apple: apply to any growth stage but application by flowering will reduce seed production

⁸Diffuse knapweed: Apply to plants in the bolting stage of development

⁹Russian knapweed: apply to plants in the spring and summer that are in the bud to flowering

¹⁰Japanese knotweed: Apply to plants 0.9-1.2 m tall

¹¹Orange hawkweed: For seasonal and 12 month control, apply to plants in the bolting stage of development.

¹²Cleavers: up to 10 cm

¹³Common lamb's quarter: Prior to budding stage

¹⁴Annual sowthistle: Up to 3 cm

¹⁵Wild chervil: Up to budding stage and 40 cm

TANK-MIX COMBINATIONS WITH MILESTONE NXT HERBICIDE

Milestone NXT Herbicide can be tank mixed with 2,4-D herbicide to broaden the spectrum of weeds controlled.

Herbicide Component	Rate	Weeds Controlled
Milestone NXT Herbicide	117 g/ha	Control of annual sow thistle, bluebur, burdock (<4 leaf), cocklebur, common plantain, flixweed, goat's beard, prickly lettuce, ragweeds, stinging nettle, sweet clover, curled dock (<4 leaf), hawkweed, peppergrass, season long control of Canada thistle, spotted knapweed, scentless chamomile, Canada goldenrod, and top growth control of blue lettuce, bull thistle, buttercup, gumweed, hoary cress, perennial sow thistle Plus all the weeds controlled by 117 g/ha of Milestone NXT alone and all the weeds listed on the 2,4-D label at 840 g ae/ha
2,4-D	840 g ae/ha	
Herbicide Component	Rate	Weeds Controlled
Milestone NXT Herbicide	150 g/ha	all weeds listed above plus season long control of absinth wormwood, dandelion
2,4-D	1080 g ae/ha	Plus all the weeds controlled by 150 g/ha of Milestone NXT alone and all the weeds listed on the 2,4-D label at 1080 g ae/ha
Herbicide Component	Rate	Weeds Controlled
Milestone NXT Herbicide	200 g/ha	all weeds listed above plus season long control of common tansy
2,4-D	1440 g ae/ha	Plus all the weeds controlled by 200 g/ha of Milestone NXT alone and all the weeds listed on the 2,4-D label at 1440 g ae/ha

Tank-Mix Combination – Milestone NXT Herbicide plus Garlon XRT Herbicide – BRUSH CONTROL

The Garlon XRT Herbicide tank-mix will provide control of a broad spectrum of broadleaf weeds, woody plants and tree species.

Use 100-200 g/ha of Milestone NXT Herbicide tank-mixed with 2.5 – 5 L/ha of Garlon XRT Herbicide. Use a surfactant such as Gateway™ Adjuvant, or non-ionic surfactant at the rate of 0.375% by volume (375 mL per 100 L of water). See surfactant label for a full list of recommended rates. See Garlon XRT Herbicide label for the full list of woody plants and trees controlled.

Tank-Mix Combination With Milestone NXT Herbicide – TOTAL VEGETATION CONTROL

The glyphosate or Arsenal Herbicide tank mix will provide control of most grass and broadleaf species.

Milestone NXT Herbicide can be tank mixed with glyphosate (present as isopropylamine salt, diammonium salt, potassium salt or dimethylamine salt, registered for use on non-cropland areas), **or** Arsenal Herbicide to broaden the spectrum of weeds, woody plants and trees controlled on rights-of-way, industrial areas (including but not limited to right-of-way and military bases) and other non-crop areas (where bare ground is desirable).

Tank-Mix Combination – Milestone NXT Herbicide plus Glyphosate Herbicide

Use 100 – 200 g/ha Milestone NXT Herbicide tank mixed with 0.8 – 4.3 kg ai/ha glyphosate herbicide (Equivalent to 1.69 – 9.0 L/ha of VP480 Herbicide). A surfactant is required. Follow mixing instructions.

Tank-Mix Combination – Milestone NXT Herbicide plus Arsenal Herbicide

Apply 2.34 - 3.0 L of Arsenal Herbicide per hectare tank mixed with 100 – 200 g/ha of Milestone NXT Herbicide.

MIXING METHODS

Mixing with Water

1. Fill the spray tank 3/4 full of clean water
2. Add the required amount of Milestone NXT Herbicide with the agitation running. Pre-slurrying with water may be necessary where there is little or no agitation or an injection system is being used or where herbicide is first added to a tank other than the spray tank.
3. If tank mixing, add the required amount of the tank-mix partner with the agitation running.
4. Add the surfactant.
5. Add antifoaming agent, if required.

SPRAYER CLEAN-OUT INSTRUCTIONS

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:
 - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
 - Agitate and circulate for 15 minutes, and flush through booms and hoses.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.
3. Second rinse:
 - Fill the tank with clean water.
 - Add All Clear Spray Tank Decontaminator, or Clean-Out Spray Tank Cleaner, or 1 L of household ammonia (containing a minimum of 3 % ammonia) per 100 L of water, or similar tank cleaning agent as per manufacturer's recommendations while filling the tank with clean water.
 - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
 - After flushing the boom and hoses, drain tank completely.
 - Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
4. Third rinse:
 - Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty odour which may cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

INVASIVE PLANT MANAGEMENT

Invasive Plant Management Practices

Milestone NXT Herbicide can be an important component of integrated vegetation management programs designed to restore desired plant communities. To maximize and extend the benefits of weed control provided by the Milestone NXT Herbicide, it is important that other vegetation management practices, including proper grazing management, fertilization, prescribed fire, mechanical mowing, hand-pulling, bio control and other methods, be used in appropriate sequences and combinations to further alleviate the adverse effects of invasive plants on desirable plant species and to promote development of desired plant communities. Consult local agriculture and land management resource specialists and/or invasive plant councils. These organizations can provide guidance on best management practices and the development of integrated vegetation management programs. Rapid response to the early detection of new invasive plants is a key invasive plant management strategy. Containment, eradication and control are essential for the management of spreading and established invasive plants.

Rangeland and Pasture Areas

For best results, invasive plants should be treated when they are actively growing and under conditions favourable for growth. Milestone NXT Herbicide provides control of emerged seedlings of susceptible plants following application.

To reduce the spread of invasive plants, it is important to treat identified invasive plants in the target area while observing required spray buffer zones for broadcast applications. Return when the wind is blowing from the opposite direction to complete spraying the targeted plant population.

Individual Plant or Spot/Strip Applications

An important component of Invasive Plant Management programs is early detection and eradication. The instructions for individual plant or spot or strip applications can be used to stop the spread of invasive plants. This is particularly important in riparian areas, where spot application is preferable to broadcast treatment.

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Aerial Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Use Milestone NXT Herbicide alone, or tank mixed with 2,4-D herbicide, glyphosate herbicide or Garlon XRT Herbicide, as a broadcast treatment by air to control listed broadleaf weeds and woody plants. Apply Milestone NXT Herbicide at the recommended rates in a minimum spray volume of 30 L/ha by air. For better coverage use 50 litres/ha. Follow all instructions under Mixing Methods for surfactant requirements and order of mixing.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate swath marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3852 or obtain technical advice from the distributor or your provincial agricultural representative.

SPRAY DRIFT PRECAUTIONS AND SPRAY BUFFER ZONES

Do not apply to any body of water. Take care to minimize the incidental overspray along the shoreline when applying to terrestrial plants at the water's edge or to water in areas where surface water is present.

Avoid spray drift at the application site. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt).

The following drift management requirement must be followed to avoid off-target drift movement from aerial applications:

1. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,

For application to rights-of-way, spray buffer zones for protection of sensitive terrestrial habitats are not required; however, the best available application strategies which minimize off-site drift, including meteorological conditions (e.g., wind direction, low wind speed) and spray equipment (e.g., coarse droplet sizes, minimizing height above canopy), should be used. Applicators must, however, observe the specified spray buffer zones for protection of sensitive aquatic habitats.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop	Aircraft type	Spray Buffer Zones (metres) Required for the Protection of:		
			Freshwater Habitat of Depths:		Terrestrial Habitat:
			Less than 1 m	Greater than 1 m	
Field sprayer	Industrial areas, other non-crop areas, rangeland and permanent pasture*	NA	5	3	10
Aerial	Industrial areas, other non-crop areas, rangeland and permanent pasture*	Fixed wing	225	150	200
		Rotary wing	100	70	150

* Spray buffer zones for the protection of terrestrial habitats are not required for use on rights-of-way including railroad ballast, rail and hydro rights-of-way, utility easements, roads, and training grounds and firing ranges on military bases.

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Milestone NXT Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Milestone NXT Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Development of plant populations resistant to this herbicide mode of action is usually not a problem on rangeland, permanent grass pastures, or non-cropland sites since these sites receive infrequent pesticide applications.
- Similar looking biotypes of a given weed species occurring in a treated area may vary in their susceptibility to a herbicide. Application of a herbicide below its labeled rate may allow more tolerant weeds to survive and a shift to more tolerant biotypes within the treated area.
- Where possible, rotate the use of Milestone NXT Herbicide or other Group 4 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Corteva Agriscience at 1-800-667-3852 or at www.corteva.ca.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

TMTrademark of Corteva Agriscience and its affiliated companies.

All other products listed are registered trademarks of their respective companies.

082024

Label Code: CN-34728-002-E

Replaces: CN-34728-001-E

Specimen Label Notes

Updated tank mix statement and address change