



# Simplicity™ GoDRI™ Herbicide

GROUP	2	HERBICIDE
-------	---	-----------

For postemergent control of wild oats, Japanese brome, yellow foxtail plus several annual broadleaf weeds including cleavers and hemp nettle in spring wheat, durum wheat, winter wheat and triticale.

## AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: pyroxsulam 21.5%  
Wettable granules

REGISTRATION NO. 31916 PEST CONTROL PRODUCTS ACT

## CAUTION: EYE IRRITANT

NET CONTENTS: 0.5 kg - bulk

**Dow AgroSciences Canada Inc.**  
2400, 215 – 2<sup>nd</sup> Street S.W.  
Calgary, Alberta  
T2P 1M4  
1-800-667-3852

**PRECAUTIONS**  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION EYE IRRITANT**

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

**At all times:** Wear clean clothing with long sleeved shirt and long pants. Wear coveralls over long sleeved shirt and long pants, chemical-resistant gloves, socks and chemical resistant footwear during mixing, loading, application, clean up and repair. Wear goggles or face shield during mixing/loading. For closed cab or aerial application, coveralls and gloves are not necessary.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

**At completion of spraying or end of the day:** Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.

**FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**TOXICOLOGICAL INFORMATION**

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

**AGRICULTURAL CHEMICAL**

Do not ship or store with food, feeds, drugs or clothing.

**ENVIRONMENTAL PRECAUTIONS**

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body. This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**STORAGE**

Keep in original container during storage. Store product in cool, dry, well ventilated place away from seed, feed, fertilizer, or other pesticides. Keep away from fire or open flame, or other source of heat. To prevent contamination, store this product away from food or feed.

## **DISPOSAL**

### **Recyclable Containers**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

### **Refillable Containers**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

### **Returnable Containers**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

### **Product**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## **GENERAL INFORMATION**

Simplicity™ GoDRI™ Herbicide is a selective postemergence herbicide for the control of wild oats and certain broadleaved weeds in spring wheat, durum wheat, winter wheat and triticale not underseeded with legumes. Simplicity GoDRI Herbicide is mixed with water and applied as a uniform broadcast spray. It is non-corrosive, nonflammable, and nonvolatile.

Simplicity GoDRI Herbicide must be applied early postemergence, to the main flush of actively growing wild oats and broadleaved weeds. Conditions favouring active weed growth enhance the activity of Simplicity GoDRI Herbicide by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details.

Simplicity GoDRI Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

### **MODE OF ACTION**

Simplicity GoDRI Herbicide inhibits the production of the ALS enzyme in plants. This enzyme is essential for the production of certain amino acids which are essential for plant growth. Simplicity GoDRI Herbicide is a Group 2 mode of action herbicide.

## **GENERAL USE PRECAUTIONS**

### **Non-Target Sites**

Do not apply where proximity of susceptible crops (e.g. canola and legumes) or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Hazards section of the label.

### **Crop Rotation**

Fields previously treated with Simplicity GoDRI Herbicide can be seeded 11 months following an application of Simplicity GoDRI Herbicide, with the following crops: barley, brown mustard, canola, dry bean (of the species *Phaseolus vulgaris*), flax, canola quality Brassica juncea, lentils, oats, field peas, chickpea, spring wheat, soybean and yellow mustard or fields can be summerfallowed. Fields previously treated with Simplicity GoDRI Herbicide can be seeded 10 months following an application of Simplicity GoDRI Herbicide with the following crops: field corn, sunflower and potatoes.

### **Tank Mixtures**

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Dow AgroSciences Canada Inc. at 1-800-667-3852 or [www.corteva.ca](http://www.corteva.ca) for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by Dow AgroSciences Canada Inc.

When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.

### **Spray Equipment Precaution**

Do not apply through any type of irrigation system.

### **Sprayer Clean-Out Instructions**

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:
  - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
  - Agitate and circulate for 15 minutes, and flush through booms and hoses.
  - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
  - Drain tank completely.
3. Second rinse:
  - Fill the tank with clean water.
  - Add All Clear Spray Tank Decontaminator as per manufacturer's recommendations while filling the tank with clean water.
  - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
  - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
  - After flushing the boom and hoses, drain tank completely.
  - Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).

4. Third rinse:
  - Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
  - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
  - Drain tank completely.

## DIRECTIONS FOR USE

**READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDED WITH LEGUMES.**

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

## SIMPLICITY GoDRI HERBICIDE ALONE

### Application Directions - Spring Wheat, Durum Wheat, Triticale

For control of low wild oat populations (<75 plants/m<sup>2</sup>) and Japanese brome from the 1 - 4 leaf (2 tiller stage, under good growing conditions), apply Simplicity GoDRI Herbicide at a rate of 52 g per hectare. Apply Simplicity GoDRI at a rate of 70 g per hectare for control of grass and broadleaf weeds (see below).

Always add a non-ionic surfactant (Agral 90, Ag-Surf or Surf 92) at 0.25 v/v (0.25 L per 100 L of spray solution) when applying Simplicity GoDRI Herbicide alone. Apply using aerial equipment or ground equipment that will assure uniform coverage. See table below for weeds species controlled. Simplicity GoDRI Herbicide will control only emerged weeds. Weeds that germinate after application will not be controlled.

### Weeds Controlled with Simplicity GoDRI at the 52 g/ha rate

oats, wild (low wild oat populations; <75 plants/m<sup>2</sup>)  
brome, Japanese (1-4 leaf) ††

### Weeds Controlled or Suppressed with Simplicity GoDRI at 70 g/ha rate

#### Weeds Controlled:

barnyard grass (1-5 leaf)	foxtail, yellow (1-5 leaf)
brome, Japanese (1-6 leaf) ††	hemp nettle (1-8 leaf)
canola, volunteer† (1-6 leaf)	oats, wild (up to 4 leaf, 2 tiller)
chickweed, common (up to 10 cm)	pigweed, red-root (1-8 leaf )

cleavers (up to 6 whorl)	round-leaved mallow (up to 6 leaf stage, <10 cm in size)
corn spurry (up to 2 whorl stage, <10 cm in height)	smartweed (lady's-thumb, 1-5 leaf)
cow cockle (up to 8 leaf)	shepherd's purse (up to 30 cm tall)
flixweed (up to 10 cm)	stinkweed (up to 30 cm tall)

† will not control volunteer Imidazolinone-tolerant canola

†† For control of Japanese brome from the 1 leaf to 4 leaf, 2 tiller stage under good growing conditions, apply Simplicity GoDRI Herbicide at the 52 g/ha rate. For larger Japanese brome (up to the 6 leaf stage) and less than optimum growing conditions, use the 70 g/ha rate.

Weeds Suppressed:

brome, downy (2-6 leaf, 4 tillers)	wild buckwheat (1-4 leaf)
dandelion (seedlings and over-wintered rosettes less than or equal to 20 cm)	Canada thistle (up to 30 cm tall, prebud)
foxtail, green (1-5 leaf)	Russian thistle (up to 10 cm tall)
white cockle (up to the first flower stage, less than 20 cm in height)	Persian darnel (1 leaf to 4 leaf, 2 tiller)

**Application Timing**

For spring wheat and durum wheat: Apply to actively growing wheat from the 3 leaf stage to before the flag leaf stage. For triticale: apply to actively growing triticale from the 3 leaf stage until the 1<sup>st</sup> node. Apply when weeds are actively growing: grass weeds up to the 4 leaf, 2 tiller stage and broadleaved weeds as indicated in the table above. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed density, control may be reduced.

**Application Directions - Winter Wheat (Fall or Spring Application) for Downy Brome**

Simplicity GoDRI Herbicide applied in the fall will provide control of downy brome while a spring application will provide suppression. Apply Simplicity GoDRI Herbicide at a rate of 70 g per hectare. Add BASF Assist Oil Concentrate at 0.8% v/v (0.8 L/100 L of spray mixture) or Merge Adjuvant at 0.5% v/v (0.5 L/100 L of spray mixture; spring application only). Apply using aerial equipment or ground equipment that will assure uniform coverage. In addition to suppression of downy brome when applied in the spring Simplicity GoDRI Herbicide will also control/suppress weeds listed. Simplicity GoDRI Herbicide will control only emerged weeds. Weeds that germinate after application will not be controlled.

**Application Timing**

Apply in the fall when winter wheat is in the 1-3 leaf stage or in the spring when winter wheat is in the 2-7 leaf, 4 tiller stage. For fall applications, apply when downy brome is actively growing at the 1 to 3 leaf stage; for spring applications, apply when weeds are actively growing at the stages indicated. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed density, control may be reduced.

Occasionally slight yellowing or height reduction may be observed in the treated crop. These transient symptoms disappear within 14 days with no reduction to yield. Do not apply to crops suffering from drought, nutrient deficiency or exposed to frost or other agronomic factors affecting plant growth.

If foliage is wet at the time of application control may be decreased. Applications of Simplicity GoDRI Herbicide are rainfast within 2 hours after application.

### Mixing Instructions (Simplicity GoDRI Herbicide used alone)

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of Simplicity GoDRI Herbicide
4. Fill the sprayer tank with sufficient water to spray 30-100 L of spray mixture per hectare.
5. Add the required surfactant
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

### Pre-harvest/Grazing Intervals

Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.

### TANK MIX COMBINATIONS

Simplicity GoDRI Herbicide can be tank mixed with herbicides and/or fungicides as indicated in the table below. When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.

**For application of Simplicity GoDRI Herbicide in tank mixtures, only add surfactant where the label of the tank mix partner product indicates its use is required, in accordance with that label. See table below for surfactant recommendations for specific tank mixtures.**

### Tank Mix Combinations with Simplicity GoDRI Herbicide at 70 g/ha

#### Spring Wheat and Durum Wheat Only

Herbicide Tank Mix Partner	Rate of herbicide TM partner (Product /ha)	Additional Pests Controlled
<b>Curtail™ M Herbicide</b>	1.5 L/ha	annual sowthistle (1-5 leaf) consult label
<b>Frontline™ XL Herbicide</b>	1.25 L/ha	annual sowthistle (1-5 leaf) stork's-bill (1-8 leaf) consult label
<b>Frontline 2,4-D XC Herbicide</b>	66.7 mL/ha Frontline 2,4-D XC A + 567 mL/ha Frontline 2,4-D XC B	annual sowthistle (1-5 leaf) stork's-bill (1-8 leaf) wild buckwheat (1-7 leaf)+ weeds controlled by 2,4-D ester at 350 g a.e./ha <sup>††</sup>

#### Spring Wheat, Durum Wheat, and Winter Wheat

Herbicide Tank Mix Partner	Fungicide Tank-Mix Partner	Rate of herbicide TM partner (Product /ha)	Additional Pests Controlled
<b>Simplicity GoDRI Herbicide alone</b>	Tilt 250 E or Stratego 250 EC	Consult fungicide tank-mix partner label for fungicide rates.	Consult fungicide tank-mix partner label for diseases controlled
<b>Attain™ XC A Herbicide + Attain XC B Herbicide</b>	Tilt 250 E or Stratego 250 EC	0.23 L/ha Attain XC A Herbicide + 0.64 L/ha Attain XC B Herbicide  Consult fungicide tank-mix partner label for fungicide rates	stork's-bill (1-8 leaf) wild buckwheat + labelled weeds for Low Rate of Attain XC A Herbicide + Attain XC B Herbicide  Consult fungicide tank-mix partner label for diseases controlled

<b>OctTain™ XL Herbicide</b>		1.1 L/ha	Consult label
<b>Spectrum™ Herbicide</b>		100 mL/ha Spectrum A + 1.5 L/ha Spectrum B	consult label
<b>Prestige™ XC Herbicide</b>		0.31 L/ha Prestige XC A + 1.5 L/ha Prestige XC B	annual sowthistle (1-5 leaf) wild buckwheat (1-9 leaf) Russian thistle (2-10 cm high)
<b>Prestige™ XL Herbicide</b>		1.75 L/ha Prestige XL	annual sowthistle (1-5 leaf) wild buckwheat (1-9 leaf) Russian thistle (2-10 cm high)
<b>Stellar™ A Herbicide + Stellar B Herbicide</b>		1.0 L/ha Stellar A Herbicide + 580 mL/ha Stellar B Herbicide	Consult label
<b>Stellar™ XL Herbicide</b>		1.0 L/ha Stellar XL Herbicide	Consult label
<b>Pixxaro™ A Herbicide</b>		308 mL/ha Pixxaro A Herbicide	Consult label
<b>Paradigm™ Herbicide</b>		25 g/ha Paradigm Herbicide	Consult label
<b>MCPA ester</b>	Tilt 250 E or Stratego 250 EC	348 – 552 g a.e./ha (e.g, 0.58 – 0.92 L/ha of MCPA Ester 600)  Consult fungicide tank-mix partner label for fungicide rates	stork's-bill (1-8 leaf) ††† wild buckwheat†+ weeds labelled for this rate range of MCPA ester  Consult fungicide tank-mix partner label for diseases controlled
		280 g a.e./ha (e.g. 465 mL/ha of MCPA Ester 600)	volunteer canola (1-6 leaf stage)* lamb's quarters (1-6 leaf stage)  *Including imidazolinone tolerant volunteer canola
<b>2,4-D ester</b>		280 g a.e./ha (e.g. 425 mL/ha of 2,4-D Ester 700)	volunteer canola (1-6 leaf stage)* lamb's quarters (1-6 leaf stage)  *Including imidazolinone tolerant volunteer canola
<b>Buctril M</b>		1.0 L/ha	consult label
<b>Refine Extra 75 DF</b>		20 g/ha A registered adjuvant is required for this mixture.	consult label



<b>Refine SG Herbicide</b>		30 g/ha A registered adjuvant is required for this mixture	consult label
<b>PP-3317 Herbicide</b>		30 g/ha A registered adjuvant is required for this mixture	consult label
<b>PP-2525 Herbicide</b>		30 g/ha A registered adjuvant is required for this mixture	consult label
<b>Thumper Emulsifiable Selective Weedkiller</b>		1.0 L/ha	consult label

†For larger weeds or heavy infestations use MCPA ester at 630 g a.e./ha (e.g., 1.05 L/ha MCPA Ester 600)

††Seedlings at the 2-4 leaf stage; actively growing mustard (except dog and green tansy), flixweed, bluebur, burdock, cocklebur, field horsetail (top growth control only), stinkweed, goats beard, prickly lettuce, hoary cress (top growth control only), kochia, lamb's-quarters, plantain, ragweeds, Russian knapweed, Russian thistle, shepherd's-purse, annual sunflower, sweet clover, vetch, wild radish

†††Use the 630 g a.e./ha rate for control of stork's-bill

**Tank Mix Combinations with Simplicity GoDRI Herbicide at 52 g/ha in Spring Wheat, Durum Wheat and Winter Wheat Under Low Wild Oat Populations (<75 plants/m<sup>2</sup>)**

<b>Herbicide Tank Mix Partner</b>	<b>Fungicide Tank Mix</b>	<b>Rate (Product /ha)</b>	<b>Additional Weeds Controlled</b>
<b>Attain XC A Herbicide + Attain XC B Herbicide</b>	Tilt 250 E or Stratego 250 EC	0.23 L/ha Attain XC A Herbicide + 0.64 L/ha Attain XC B Herbicide  Consult fungicide tank-mix partner label for fungicide rates.	cow cockle (up to 8 leaf), wild oats, wild buckwheat (1-4 leaf) + labelled weeds for Low Rate of Attain XC A Herbicide + Attain XC B Herbicide  Consult fungicide tank-mix partner label for diseases controlled
<b>OcTTain™ XL Herbicide</b>		1.1 L/ha	Consult label
<b>Stellar A Herbicide + Stellar B Herbicide</b>		1.0 L/ha Stellar A Herbicide + 580 mL/ha Stellar B Herbicide	Consult label
<b>Stellar XL Herbicide</b>		0.8 - 1.0 L/ha	Consult label

**Mixing Instructions (Simplicity GoDRI Herbicide tank mixed with other products)**

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of Simplicity GoDRI Herbicide
4. Add the correct amount of tank-mix partner and continue to agitate.
5. Fill the sprayer tank with the remaining amount of water required.

**Notes: Follow tank-mix partner label for order of mixing.**

## **APPLICATION INSTRUCTIONS (GROUND AND AERIAL APPLICATIONS)**

### **(1) Ground Application (Simplicity GoDRI Herbicide applied alone and with all tank mix partners)**

Using ground equipment, apply Simplicity GoDRI Herbicide alone or with tank mix partners as a broadcast treatment. Apply Simplicity GoDRI Herbicide alone at the recommended rates in a spray volume of 50-100 L water / ha. When tank mixing refer to the label of the tank mix partner label for additional instructions, water volumes, precautions and weeds controlled.

### **(2) Aerial Application**

#### **Directions for Use**

**Do not use human flaggers.**

Use Simplicity GoDRI Herbicide alone or in tank mix with either MCPA ester or Buctril M as a broadcast treatment by air. Apply the recommended rates of Simplicity GoDRI Herbicide and either MCPA ester or Buctril M as indicated above. Apply in a minimum spray volume of 30 L water / ha. Refer to the tank mix partner label for additional instructions, precautions and weeds controlled.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate swath marking devices.

#### **Use Precautions**

Apply only when meteorological conditions at the treatment site allow for complete and even coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

#### **Operator Precautions**

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing should be laundered regularly.

#### **Product Specific Precautions**

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3852 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

- The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.
- The following drift management requirement must be followed to avoid off-target drift movement from aerial applications:  
Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

## BUFFER ZONES

Spot treatments using hand-held equipment do not require a buffer zone.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:	
			Freshwater Habitat	Terrestrial Habitat
Field sprayer	Spring wheat, durum wheat, winter wheat, triticale		1	2
Aerial	Spring wheat, durum wheat, winter wheat, triticale	Fixed wing	1	70
		Rotary wing	1	60

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

NOTE: Applicators may recalculate a site-specific buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. To access the Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

## RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Simplicity GoDRI Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Simplicity GoDRI Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Simplicity GoDRI Herbicide or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Dow AgroSciences Canada Inc. at 1-800-667-3852 or at [www.corteva.ca](http://www.corteva.ca).

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

<sup>TM</sup>Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.

All products listed are registered trademarks of their respective companies.

011620

Label Code: CN-31916-011-E

Replaces: CN-31916-010-E

Label Change : Addition of triticale to label