

Acapela™

Speed, agility and exceptional coverage.

FUNGICIDE

WHY USE ACAPELA™ FUNGICIDE?

- Acapela™ fungicide is rapidly absorbed, moving quickly into and within each plant allowing you to spray even when conditions are challenging.
- Acapela is a one-of-a-kind fungicide that quickly and efficiently surrounds, penetrates, and protects the leaf and stem.
- Acapela is a Group 11 fungicide that controls sclerotinia in canola.
- In addition to outstanding disease control, Acapela supports positive plant performance, even in stressful conditions by increasing chlorophyll content and plant productivity.
- Better coverage means more consistent protection, providing outstanding disease control for greener, healthier crops and higher yield potential.

DISEASES CONTROLLED

Canola

- Sclerotinia rot – white mould

Cereals

- Crown rust
- Leaf rust
- Net blotch
- Powdery mildew
- Scald
- Septoria leaf blotch
- Stripe rust
- Tan spot

Corn

- Northern corn leaf blight

Flax

- Pasmio (*Septoria linicola*)

Pulse crops (peas, lentils, chickpeas, dry beans)

- Anthracnose (*lentils and dry beans*)
- Ascochyta blight (*in lentils*)
- Asian soybean rust
- Mycosphaerella blight* (*field peas*)
- Sclerotinia rot – white mould*

Soybeans

- Asian soybean rust
- Frogeye leafspot
- Sclerotinia rot – white mould*
- Septoria brown spot

Potatoes

- Early blight
- White mould (*Sclerotinia sclerotiorum*)
- Late blight

IMPORTANT PRECAUTIONS

1. Do not apply to cereals after flowering (Feekes 10.5.1 or Zadoks 60).
2. Maximum seasonal use rate is 2.64 L/ha.

APPLICATION GUIDELINES

<p>CROPS</p> <p>Canola</p> <p>Cereals</p> <p>Corn</p> <p>Flax</p> <p>Potatoes</p> <p>Pulse crops (peas, lentils, chickpeas, dry beans)</p> <p>Soybeans</p>	<p>PACKAGING</p> <ul style="list-style-type: none"> • 9.6 L jug • 115.2 L drum
	<p>RATES & ACRES TREATED</p> <ul style="list-style-type: none"> • Canola: 0.32 L/ac (30 ac/jug or 360 ac/drum) • Cereals: 0.19 L/ac (50 ac/jug or 600 ac/drum) • Corn: 0.21 to 0.32 L/ac (30 to 45 ac/jug) • Flax: Pasmó (<i>Septoria linicola</i>) 0.24 L/ac (40 ac/jug or 480 ac/drum) • Pulse crops: (Peas, lentils, chickpeas, dry beans) <ul style="list-style-type: none"> - Anthracnose, Ascochyta blight: 0.24 L/ac (40 ac/jug or 480 ac/drum) - Sclerotinia rot - white mould: 0.35 L/ac (27 ac/jug or 325 ac/drum) • Soybeans: <ul style="list-style-type: none"> - Sclerotinia rot - white mould: 0.35 L/ac (27 ac/jug) - Asian soybean rust, frog-eye leafspot, septoria brown spot: 0.24 to 0.35 L/ac (27 to 40 ac/jug) • Potatoes: <ul style="list-style-type: none"> - Early blight and white mold: 0.24 - 0.40 L/ac - Late blight: 0.18 - 0.40 L/ac
	<p>WATER VOLUME</p> <ul style="list-style-type: none"> • Ground minimum of 110 L/ha (10 US gal/ac) • Aerial minimum of 50 L/ha (4.5 US gal/ac)
	<p>TIMING</p> <p>Refer to the Acapela label for complete use instructions.</p>
	<p>RAINFAST</p> <p>1 hour</p>

MIXING INSTRUCTIONS

1. Shake well before use
2. Fill clean spray tank $\frac{1}{3}$ to $\frac{1}{2}$ full of water
3. While agitating, add the required amount of Acapela fungicide, continuing agitation until the product is completely dispersed
4. Continue filling the tank with agitation. Mix thoroughly to fully disperse the fungicide; once dispersed continued agitation is required. Use mechanical or hydraulic means; do not use air agitation



CROP ROTATION

Any crop the following year.