

Amity™ WDG

HERBICIDE

Superior control of grass and broadleaf weeds in Clearfield® canola

WHY USE AMITY™ WDG HERBICIDE?

- Provides broad spectrum weed control for Clearfield canola.
- Provides rotational flexibility.
- Reliable control of tough grasses and broadleaf weeds.
- Tank mix with Lontrel™ XC herbicide for improved wild buckwheat and Canada thistle control.

BROADLEAF WEEDS CONTROLLED

- Cow cockle
- Lamb's-quarters
- Redroot pigweed
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Wild mustard
- Volunteer canola (excluding Clearfield® varieties)

GRASS WEEDS CONTROLLED

- Barnyard grass
- Green foxtail
- Persian dandelion
- Volunteer barley
- Volunteer canaryseed
- Volunteer durum wheat
- Volunteer spring wheat (excluding Clearfield® varieties)
- Volunteer tame oats
- Wild oats
- Yellow foxtail

WEEDS SUPPRESSED

- Cleavers
- Japanese brome
- Round-leaved mallow
- Wild buckwheat

Surjet™

ADJUVANT

Surjet™ adjuvant is a blend of surfactant and petroleum hydrocarbons designed for use with Ares™ SN or Amity WDG herbicides.

Rate: 0.5% v/v

Water volume: 40 L/ac (10 US gal/ac)

Surjet adjuvant sold separately from Amity™ WDG herbicide.

APPLICATION GUIDELINES

CROPS Clearfield® Canola	PACKAGING	4 x cartons with 4 x pouches of 117.5g wettable dry granule
	RATES & ACRES TREATED	<ul style="list-style-type: none"> • 10 ac/pouch • 40 ac/carton • 160 ac/case
	WATER VOLUME	<ul style="list-style-type: none"> • Ground 40 L/ac (10 US gal/ac) • Add Surjet™ adjuvant at 0.5% v/v • Aerial NOT registered
	TIMING	Crop stage: 2 to 6-leaf stage Weed stage: 1 to 4-leaf stage for grassy weeds and cotyledon to 4-leaf stage for broadleaf weeds
	RAINFAST	3 hours
	TANK-MIX HERBICIDES	Lontrel™ XC herbicide



DOWNLOAD THE 2025 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.

CROP ROTATION

- Winter wheat can be planted 3 months after treatment as a rotational crop.
- Initial crop injury to non Clearfield® canola may be observed.
- Avoid spray overlap as yield reduction may result.

The following crops may be grown safely the year following an application:

- | | |
|--------------------------|--------------------------|
| • Canaryseed | • Non-Clearfield® canola |
| • Chickpeas | • Oats |
| • Clearfield® Canola | • Spring barley |
| • Clearfield® Sunflowers | • Spring wheat |
| • Durum wheat | • Sunflower |
| • Field corn | |
| • Field peas | |
| • Flax | |
| • Lentils | |

If drought conditions exist in the first season after application, do not grow non Clearfield canola, durum, flax or sunflowers.



GRAZING AND HARVEST

Do not harvest treated crop within 60 days after application.