

Amity™ WDG

HERBICIDE

Superior control of grass and broadleaf weeds in Clearfield® canola

WHY USE AMITY™ WDG HERBICIDE?

- Consistent and reliable post-emergent weed control including tough grassy and targeted broadleaf weeds
- Broad spectrum control for Clearfield canola
- Rotational flexibility for crop planning on your farm
- Tank-mix with Lontrel™ XC herbicide for improved wild buckwheat and Canada thistle control

GRASS WEEDS CONTROLLED

- Barnyard grass
- Japanese brome*
- Yellow foxtail
- Green foxtail
- Persian darnel
- Wild oats
- Volunteer canary seed
- Volunteer barley
- Volunteer tame oats
- Volunteer spring wheat
(excluding Clearfield® varieties)
- Volunteer durum wheat

BROADLEAF WEEDS CONTROLLED

- Cleavers*
- Cow cockle
- Lamb's-quarters
- Redroot pigweed
- Round-leaved mallow*
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Wild buckwheat*
- Wild mustard
- Volunteer canola
(excluding Clearfield® varieties)

Surjet™

ADJUVANT

Surjet™ Adjuvant is a blend of surfactant and petroleum hydrocarbons designed for use with Ares™ SN or Amity WDG herbicides.

Rate: 0.5% v/v

Water volume: 40 L/ac (10 US gal/ac)

Surjet Adjuvant sold separately from Amity™ WDG herbicide.

* Suppression only

APPLICATION GUIDELINES

<p>CROPS</p> <p>Clearfield® Canola</p>	<p>PACKAGING</p> <p>4 x cartons with 4 x pouches of 117.5g wettable dry granule</p>
	<p>RATES & ACRES TREATED</p> <ul style="list-style-type: none"> • 10 ac/pouch • 40 ac/carton • 160 ac/case
	<p>WATER VOLUME</p> <ul style="list-style-type: none"> • Ground 40 L/ac (10 US gal/ac) • Add Surjet at 0.5% v/v • Aerial NOT registered
	<p>TIMING</p> <p>Crop stage: 2 to 6-leaf stage Weed stage: 1 to 4-leaf stage for grassy weeds and cotyledon to 4-leaf stage for broadleaf weeds</p>
	<p>RAINFAST</p> <p>3 hours</p>
	<p>TANK MIX HERBICIDES</p> <p>Lontrel™ XC</p>

MIXING INSTRUCTIONS

1. Fill the spray tank with ½ to ¾ of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add required amount of Amity WDG
4. Add any tank mix partners
5. Add the required amount of Surjet™ adjuvant at 0.5% v/v
6. Complete filling the sprayer tank with water

Note: Do not mix undiluted herbicides in the chem-handler

CROP ROTATION

- Winter wheat can be planted 3 months after treatment as a rotational crop.
- Initial crop injury to non-Clearfield canola may be observed.
- Avoid spray overlap as yield reduction may result.

The following crops may be grown safely the year following an application:

- Canary seed
- Chickpeas
- Clearfield® Canola (canola hybrids with the Clearfield trait)
- Clearfield Sunflowers (sunflower varieties with the Clearfield trait)
- Durum wheat
- Field corn
- Field peas
- Flax
- Lentils
- Non-Clearfield canola
- Spring barley
- Spring wheat
- Sunflower

If drought conditions exist in the first season after application, do not grow non-Clearfield canola, durum, flax or sunflowers.

The following crop may be grown safely two years following an application:

- Mustard (condiment type only)
- Oats

For further information, please consult the Amity WDG label

GRAZING AND HARVEST

Do not harvest treated crop within 60 days after application.