

Ares™ SN

The performance standard in weed control for Clearfield® canola.

HERBICIDE**WHY USE ARES™ SN HERBICIDE?**

- **Consistent and reliable post-emergent weed control**, including subsequent flushes.
- **Flexible application timing** with a wide window of application on both crop and weeds.
- **Superior control of lamb's quarters, wild buckwheat, cleavers and volunteer canola.**

BROADLEAF WEEDS CONTROLLED

- Chickweed
- Cleavers^{2,3}
- Cow cockle
- Hemp-nettle²
- Lamb's-quarters²
- Redroot pigweed
- Round-leaved mallow

- Russian thistle
- Shepherd's purse²
- Smartweed²
- Stinkweed
- Stork's-bill
- Wild buckwheat
- Wild mustard
- Volunteer tame mustard
- Volunteer canola (excluding Clearfield® varieties)

GRASS WEEDS CONTROLLED

- Barnyard grass
- Japanese brome¹
- Yellow foxtail
- Green foxtail
- Persian dandelion
- Wild oats
- Volunteer canaryseed
- Volunteer barley

- Volunteer tame oats
- Volunteer spring wheat (excluding Clearfield® varieties)
- Volunteer durum wheat

Surjet™**ADJUVANT**

Surjet™ adjuvant is a blend of surfactant and petroleum hydrocarbons designed for use with Ares™ SN or Amity WDG herbicides.

Rate: 0.5% v/v

Water volume: 40 L/ac (10 US gal/ac)

Surjet adjuvant is sold in a co-pack with Ares SN.

¹ For spring germinating Japanese brome control apply at 1 to 4-leaf stage.
² Group 2 susceptible biotypes will not be controlled.

³ A tank mix with Facet® L will improve control of Group 2 resistant biotypes.

APPLICATION GUIDELINES

CROPS Clearfield® Canola	PACKAGING	Case: <ul style="list-style-type: none"> • Ares™ SN herbicide: 1 x 98 L jug • Surjet™ adjuvant: 1 x 8.1 L jug
	RATES & ACRES TREATED	Rate: 244 mL/ac Acres: 40 ac/case
	WATER VOLUME	<ul style="list-style-type: none"> • Ground 40 L/ac (10 US gal/ac) • Aerial NOT registered
	TIMING	Crop stage: 2 to 7-leaf stage Weed stage: 1 to 6-leaf stage for grassy weeds and cotyledon to 4-leaf stage for broadleaf weeds
	RAINFAST	2 hours
	TANK-MIX HERBICIDES	Lontrel™ XC herbicide Facet® L herbicide



DOWNLOAD THE 2025 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.

CROP ROTATION

Sensitivity of injury can vary depending on the crop. If rainfall in your area was less than 125 mm during the growing season between **June 1 to August 31**, it is recommended that you grow field peas, Clearfield® lentils or Clearfield® wheat on Clearfield® canola stubble.

The following crops may be grown safely the year following an application:

- Canaryseed
- Chickpeas
- Field peas
- Field corn
- Clearfield® canola
- Clearfield® canola-quality Brassica juncea
- Lentils (including Clearfield® lentils)
- Spring wheat (including Clearfield® varieties)
- Spring barley

The following crops may be grown safely two years following an application:

- Canola
- Flax
- Durum wheat
- Sunflower

GRAZING AND HARVEST

Do not harvest treated crop within 60 days after application.

