

Ares™ SN

The performance standard in weed control for Clearfield® canola.

HERBICIDE

WHY USE ARES™ SN HERBICIDE?

- Consistent and reliable post-emergent weed control, including subsequent flushes.
- Wide window of application on both crop and weeds.
- Superior control of lamb's quarters, wild buckwheat, cleavers and volunteer canola.

GRASS WEEDS CONTROLLED

- Barnyard grass
- Japanese brome¹
- Yellow foxtail
- Green foxtail
- Persian darnel
- Wild oats
- Volunteer canary seed
- Volunteer barley
- Volunteer tame oats
- Volunteer spring wheat (excluding Clearfield® varieties)
- Volunteer durum wheat

BROADLEAF WEEDS CONTROLLED

- Chickweed
- Cleavers^{2,3}
- Cow cockle
- Hemp-nettle²
- Lamb's-quarters²
- Redroot pigweed
- Round-leaved mallow
- Russian thistle
- Shepherd's purse²
- Smartweed²
- Stinkweed
- Stork's bill

- Wild buckwheat
- Wild mustard
- Volunteer tame mustard
- Volunteer canola (excluding Clearfield® varieties)

Surjet™

ADJUVANT

Surjet™ Adjuvant is a blend of surfactant and petroleum hydrocarbons designed for use with Ares™ SN or Amity WDG herbicides.

Rate: 0.5% v/v

Water volume: 40 L/ac (10 US gal/ac)

Surjet Adjuvant is sold in a co-pack with Ares SN.

¹For spring germinating Japanese brome control apply at 1 to 4-leaf stage

²Group 2 susceptible biotypes will not be controlled

³A tank mix with Facet® L will improve control of Group 2 resistant biotypes

APPLICATION GUIDELINES

<p>CROPS</p> <p>Clearfield® Canola</p>	<p>PACKAGING</p> <p>Case:</p> <ul style="list-style-type: none"> • Ares™ SN: 1 x 9.8 L jug • Surjet Adjuvant: 1 x 8.1 L jug
	<p>RATES & ACRES TREATED</p> <p>40 ac/case</p>
	<p>WATER VOLUME</p> <ul style="list-style-type: none"> • Ground 40 L/ac (10 US gal/ac) • Aerial NOT registered
	<p>TIMING</p> <p>Crop stage: 2 to 7-leaf stage Weed stage: 1 to 6-leaf stage for grassy weeds and cotyledon to 4-leaf stage for broadleaf weeds</p>
	<p>RAINFAST</p> <p>2 hours</p>
	<p>TANK MIX HERBICIDES</p> <p>Lontrel™ XC</p>

MIXING INSTRUCTIONS

1. Fill the spray tank with ½ to ¾ of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add required amount of Ares SN
4. Add any tank mix partners
5. Add the required amount of Surjet adjuvant at 0.5% v/v
6. Complete filling the sprayer tank with water

Note: Do not mix undiluted herbicides in the chem-handler

CROP ROTATION

Sensitivity of injury can vary depending on the crop. If rainfall in your area was less than 125 mm during the growing season between **June 1 to August 31**, it is recommended that you grow field peas, Clearfield lentils or Clearfield wheat on Clearfield canola stubble.

The following crops may be grown safely the year following an application:

- Canary seed
- Chickpeas
- Field peas
- Field corn
- Clearfield canola
- Clearfield canola quality Brassica juncea
- Lentils including Clearfield lentils
- Spring wheat including Clearfield spring wheat
- Spring barley

The following crops may be grown safely two years following an application:

- Canola
- Durum wheat
- Flax
- Sunflower

GRAZING AND HARVEST

Do not harvest treated crop within 60 days after application.