

# Attain™ XC

## HERBICIDE

Extra-concentrated excellence on 40 of today's toughest broadleaf weeds in all soil zones in Western Canada.

### WHY USE ATTAIN™ XC HERBICIDE?

- Weed control. Gets tough weeds like kochia, stork's-bill, round-leaved mallow and more.
- Choice. Choose the rate option that suits your weed pressure – 40 or 53 ac/case.

#### 40 AC/CASE RATE

##### WEEDS CONTROLLED

- Blue lettuce<sup>1</sup>
- Bluebur
- Burdock
- Cleavers<sup>3</sup>
- Cocklebur
- Dandelion<sup>2</sup>
- Docks
- Dog mustard
- Field bindweed<sup>1</sup>
- Field horsetail<sup>1</sup>
- Field peppergrass
- Flixweed
- Goat's-beard

- Gumweed
- Hairy galinsoga
- Hedge bindweed
- Hoary cress<sup>1</sup>
- Kochia<sup>3</sup>
- Lamb's-quarters
- Leafy spurge<sup>1</sup>
- Oak-leaved goosefoot
- Plantain
- Prickly lettuce
- Ragweed
- Redroot pigweed

- Round-leaved mallow
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Stork's-bill
- Sweet clover
- Tansy mustard
- Tartary buckwheat
- Vetch
- Volunteer canola (all herbicide tolerant varieties)

- Volunteer flax
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild radish

##### WEEDS SUPPRESSED

- Annual sow thistle
- Canada thistle<sup>1</sup>
- Chickweed<sup>3</sup>
- Hemp-nettle
- Perennial sow thistle<sup>1</sup>

#### 53 AC/CASE RATE

##### WEEDS CONTROLLED

- Annual sunflower
- Bluebur
- Burdock
- Cleavers<sup>3</sup>
- Cocklebur
- Field horsetail<sup>1</sup>
- Flixweed

- Goat's-beard
- Hoary cress<sup>1</sup>
- Lamb's-quarters
- Kochia<sup>3</sup>
- Plantain
- Prickly lettuce
- Ragweed

- Shepherd's purse
- Stinkweed
- Sweet clover
- Vetch
- Volunteer canola (all herbicide tolerant varieties)
- Wild buckwheat

- Wild mustard
- Wild radish

<sup>1</sup> Top growth control only

<sup>2</sup> Spring rosettes

<sup>3</sup> Including Group 2 resistant biotypes

FLUROXYPYR: 43 g/ac (Low Rate: 32 g/ac)  
2, 4-D ESTER: 227 g/ac (Low Rate: 170 g/ac)

## APPLICATION GUIDELINES

|  |   |
|--|---|
| <p><b>CROPS</b></p> <p>Barley</p> <p>Durum wheat</p> <p>Forage grasses</p> <p>Spring wheat</p> <p>Winter wheat</p> | <p><b>PACKAGING</b></p> <ul style="list-style-type: none"> <li>• Attain XC A: 1 x 5 L</li> <li>• Attain XC B: 2 x 6.8 L</li> </ul>  |
|  | <p><b>RATES</b></p> <p><b>Low rate:</b></p> <ul style="list-style-type: none"> <li>• Attain XC A: 96 ml/ac</li> <li>• Attain XC B: 260 ml/ac</li> </ul> <p><b>High rate:</b></p> <ul style="list-style-type: none"> <li>• Attain XC A: 130 ml/ac</li> <li>• Attain XC B: 340 ml/ac</li> </ul> |
|  | <p><b>ACRES TREATED</b></p> <p><b>Low rate:</b> 53 ac/case</p> <p><b>High rate:</b> 40 ac/case</p>  |
|  | <p><b>WATER VOLUME</b></p> <ul style="list-style-type: none"> <li>• Ground 20-40 L/ac (5-10 US gal/ac)</li> <li>• Aerial 12-20 L/ac (3-5 US gal/ac)</li> </ul>  |
|  | <p><b>TIMING</b></p> <p><b>Crop stage:</b> 4-leaf to just prior to flag leaf emergence</p> <p><b>Weed stage:</b> 1 to 6-leaf or as indicated on the product label</p>   |
|  | <p><b>RAINFAST</b></p> <p>1 hour</p>  |
|  | <p><b>TANK MIX HERBICIDES</b></p> <ul style="list-style-type: none"> <li>• Simplicity™ GoDRI™</li> <li>• Liquid Achieve™ SC</li> <li>• Assert®</li> <li>• Everest®</li> <li>• Horizon®</li> <li>• Puma®</li> </ul>  |

### MIXING INSTRUCTIONS

1. Fill the spray tank with ½ to ¾ of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add any required water conditioners
4. Add any tank mix partners that are a dry formulation
5. Add the required amount of grassy weed tank mix partner
6. Add the required amount of Attain XC A
7. Add the required amount of Attain XC B
8. Add any required adjuvant or surfactants
9. Complete filling the sprayer tank with water

*Note: Do not mix undiluted herbicides in the chem-handler*

### CROP ROTATION

All major crops may be grown the following spring.

### GRAZING AND HARVEST

- Allow 7 days after application before grazing lactating animals.
- Livestock may graze treated areas 3 days after application.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 30 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.