# **Delegate**<sup>\*\*</sup>

Jemvelva<sup>®</sup>active

# **INSECTICIDE**

Delegate<sup>™</sup> insecticide with Jemvelva<sup>™</sup> active (spinetoram) offers quick knockdown and residual control of Western bean cutworm and European corn borer in corn with a unique Group 5 mode of action.

## **DELEGATE OFFERS**

- Fast and effective control of Western bean cutworm and European corn borer by either contact or ingestion.
- Important Integrated Pest Management (IPM) tool with its unique mode of action Jemvelva active.
- Low use rates.

Translaminar activity provides:

- Extra protection against insects that feed from the underside of leaves
- Long-lasting residual control

# WESTERN BEAN CUTWORM CROP IMPACT

In corn, young Western bean cutworm larvae feed on the tassels and silks until they are large enough to tunnel into the ear and feed extensively on the kernels, causing yield damage and grain quality concerns.

# **IPM COMPATIBLE**

Delegate has a favourable environmental profile and has minimal effects on beneficial insects. Industry field trials have shown that Delegate has a low impact on beneficial populations of key arthropod natural enemies, such as ladybugs

#### **RESISTANCE MANAGEMENT**

Delegate should be used as part of an overall Insect Resistance Management (IRM) strategy which includes monitoring, scouting and thresholds. Alternating different insecticide groups is a preferred approach to IRM. The unique Group 5 chemistry of Delegate provides an ideal rotational mode of action for managing Western bean cutworm.

#### TANK MIXING

Delegate can be tank mixed with a fungicide, like Acapela<sup>™</sup>, provided:

- The tank mix partner is registered in the crop
- The application timing of all tank mix partners are compatible with crop and pest staging
- The most restrictive label directions of the tank mix are followed

## **OPTIMIZING PERFORMANCE**

- Rainfast when spray solution has dried on the vegetation (two hours).
- A spray solution pH between 5 and 9 is preferred for optimal performance.

5

- Ground application: Use spray equipment capable of thorough coverage of the crop, ensuring uniform coverage of the target pest.
- Aerial application: Apply in a solution of at least 30 litres per hectare.

## DELEGATE: UNIQUE MODE OF ACTION

Jemvelva<sup>™</sup> active, the active ingredient in Delegate, is the only Group 5 insecticide registered for Western bean cutworm control in corn. No other class of products affects the target pest with the same mode of action as Jemvelva active.

#### PRECAUTIONS

Delegate residues that have dried on the vegetation are non-toxic to bees. Do not spray when bees are active in the area.



Aerial applications do not require a permit.



APPLICATION GUIDELINES	
Active ingredient:	Spinetoram
Chemical group:	Group 5
Formulation:	Wettable granule
Packaging:	6 x 840 g bottles
Application	Aerial and ground application
Re-entry period:	12 hours
Pre-harvest interval:	Field and popcorn: 28 days Seed and sweet corn: 1 day
Rates:	50 to 85 g/ac (120 to 210 g/ha)
Guidelines:	Ensure sufficient water volume for complete coverage of the plant foliage
Applications should be timed at egg hatch or to small larvae. Use the higher rate for heavy infestations, large larvae or longer residual control.	
Repeat applications based on monitoring of insect populations. Apply a maximum of three applications per year with a minimum of five days between applications.	



