

Delegate™

Jemvelva™ active

Harness nature's strength for sustainable farming.

INSECTICIDE

WHY USE DELEGATE™ INSECTICIDE WITH JEMVELVA™ ACTIVE?

- **Performance.** Delegate™ provides quick and effective control of foliage feeding insects including Colorado potato beetle and European corn borer.
- **Resistance management.** Delegate contains a unique Group 5 active ingredient, Jemvelva active, making it an excellent tool for potato growers managing Colorado potato beetle resistance.
- **Ease of use.** Low use rates delivered through a convenient dry formulation.
- **Flexibility.** Market access approvals allow freedom to use Delegate across all production acres.

INSECTS CONTROLLED

CORN

- European corn borer
- Western bean cutworm

POTATOES

- Colorado potato beetle
- European corn borer

SOYBEANS


- Armyworm

WHEAT

- Armyworm

DELEGATE IS REGISTERED FOR AERIAL APPLICATION ON POTATOES

SPEED OF KNOCKDOWN OF COLORADO POTATO BEETLE 2 DAYS AFTER APPLICATION



APPLICATION GUIDELINES

CROPS Corn Potatoes Soybeans Wheat	PACKAGING 6 x 840 g bottles
	RATES <ul style="list-style-type: none"> • Corn: <ul style="list-style-type: none"> - Western bean cutworm: 50-85 g/ac - European corn borer: 50-85 g/ac • Potatoes: <ul style="list-style-type: none"> - Colorado potato beetle: 65-97 g/ac - European corn borer: 65 g/ac • Soybeans & Wheat: <ul style="list-style-type: none"> - Armyworm: 40-80 g/ac
	TIMING <ul style="list-style-type: none"> • Western bean cutworm and European corn borer: Time the application at egg hatch or to small larvae. • Armyworm: Time the application at peak egg hatch and/or small larvae stage. • Colorado potato beetle: Time the application at egg hatch or small larvae. • Use the higher rate for higher pest pressure or for larger larvae. Ensure sufficient water volume for complete coverage of the plant foliage.
	RAINFAST 2 hours
	TANK MIXES Delegate™ insecticide with Jemvelva™ active can be mixed with fungicides and micro-nutrients. Consult your Corteva Agriscience representative if you are tank-mixing Delegate™.

OPTIMIZING PERFORMANCE

- Spray solution pH can affect the performance of Delegate
 - A spray pH between 5 and 9 is preferred for best results
 - If required, adjust spray solution pH prior to the addition of Delegate
- **Aerial application (for potatoes and corn only):** Apply only by fixed-wing or rotary aircraft equipment. Use a minimum spray volume of 12 L/ac.

CROP ROTATION

Rotate to labelled crops only.

PRECAUTIONS

- Maximum of 3 applications per year
- Minimum re-treatment interval of 5 days for corn, soybeans and wheat
- Minimum re-treatment interval of 7 days for potatoes
- **Do not** apply this product to flowering crops or weeds if bees are visiting the treatment area.
- **Do not** make more than two consecutive applications of Group 5 insecticides.

PRE-HARVEST INTERVAL

- **Sweet and seed corn:** 1 day
- **Forage and potatoes:** 7 days
- **Wheat:** 21 days
- **Soybeans, field corn, popcorn and stover harvest:** 28 days

Always read and follow label directions.