

Eclipse™ XC

HERBICIDE

Broad-spectrum weed control in glyphosate-tolerant corn & canola for superior control of wild buckwheat and Canada thistle.

WHY USE ECLIPSE™ XC HERBICIDE?

- One pass solution that offers superior control of a broad spectrum of weeds.
- Resistance Management – two unique modes of action.
- Crop safety – protects the crop without negatively impacting yield or maturity.
- Convenient packaging – one case contains all you need for 40 acres.

BROADLEAF WEEDS CONTROLLED

- Chickweed
- Cleavers
- Corn spurry
- Cow cockle
- Hemp-nettle
- Kochia
- Lamb's-quarters
- Night-flowering catchfly
- Redroot pigweed

- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Volunteer canola (except glyphosate resistant biotypes)
- Wild buckwheat*
- Wild mustard
- Wild tomato

PERENNIAL WEEDS CONTROLLED

- Canada thistle¹
- Dandelion¹ (<15 cm diameter)
- Perennial sow thistle¹

GRASS WEEDS CONTROLLED

- Green foxtail
- Quackgrass¹
- Volunteer cereals
- Wild oats

WEEDS SUPPRESSED

- Dandelion (>15 cm diameter)

WILD BUCKWHEAT CONTROL

Control up to the 6-leaf stage. Glyphosate alone does not provide consistent control of wild buckwheat past the 2 to 3-leaf stage.

CANADA THISTLE CONTROL

Season-long control with the combination of Group 4 and 9 active ingredients. Experience the same control on perennial sow thistle.

APPLICATION GUIDELINES

CROPS Glyphosate tolerant canola Glyphosate tolerant corn	PACKAGING	• Eclipse XC A: 1 x 2.67 L • Eclipse XC B: 2 x 7.5 L
	RATES & ACRES TREATED	40 ac/case
	WATER VOLUME	• Ground 40 L/ac (10 US gal/ac) • Aerial not registered
	TIMING	Canola crop stage: 2 to 6-leaf Corn crop stage: VE-V6 (8-leaf stage)
	RAINFAST	4 hours
	TANK MIX HERBICIDES	None registered

MIXING INSTRUCTIONS

1. Fill the spray tank with ½ to ¾ of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add the required amount of Eclipse XC A
4. Add the required amount of Eclipse XC B
5. Add any required adjuvant or surfactants
6. Complete filling the sprayer tank with water

Note: Do not mix undiluted herbicides in the chem-handler

CROP ROTATION

The year following application, fields can be seeded to:

- Barley
- Canola
- Flax
- Forage grasses
- Mustard
- Oats
- Peas²
- Rye (not under-seeded with legumes)
- Soybeans²
- Summerfallow
- Wheat

GRAZING AND HARVEST

- For field corn, allow 40 days before grazing treated areas or feeding cattle with corn silage from treated areas.
- For canola, allow 3 to 5 days before grazing treated areas.

*Weed controlled through multi-effective mode of action

¹ Season-long control

² For field pea or soybean rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140 mm (5.5 inches) and annual rainfall must be greater than 175 mm (6.9 inches).