

Eclipse™ XC

HERBICIDE

Broad-spectrum weed control in glyphosate-tolerant corn & canola for superior control of wild buckwheat and Canada thistle

WHY USE ECLIPSE XC HERBICIDE?

- Convenient all-in-one herbicide solution that offers superior control of a broad spectrum of weeds.
- Two modes of action for resistance management.
- Protects canola without negatively impacting yield or maturity, unlike elevated rates of glyphosate.
- Stronger XC formulation. A convenient, higher concentrated solution.

WEEDS CONTROLLED

- Canada thistle
- Canola (except Roundup Ready)
- Chickweed
- Cleavers
- Corn spurry
- Cow cockle
- Dandelion (<15 cm diameter)
- Green foxtail
- Hemp-nettle
- Kochia
- Lady's-thumb
- Lamb's-quarters
- Night-flowering catchfly
- Perennial sow thistle
- Quackgrass
- Redroot pigweed
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Volunteer cereals
- Wild buckwheat*
- Wild mustard
- Wild oats
- Wild tomato

WILD BUCKWHEAT CONTROL

Control even up to the 6-leaf stage. Glyphosate alone does not provide consistent control of wild buckwheat past the 2- to 3-leaf stage.

CANADA THISTLE CONTROL

Season-long control with the combination of Group 4 and 9 active ingredients. Experience the same control on perennial sow thistle.

*Weed controlled through multi-mode of action and †Season long control

ECLIPSE XC APPLICATION GUIDELINES

CROPS: Glyphosate tolerant corn and canola	RATE AND PACKAGING <ul style="list-style-type: none"> • 40 ac/case • 240 ac/drum and case • 960 ac/bulk pallet • Water volume <ul style="list-style-type: none"> • Ground 40 L/ac (10 US gal/ac) • Aerial not registered
	WHEN TO APPLY: <ul style="list-style-type: none"> • Corn stage: VE - V6 • Canola stage: 2 - 6 leaf
	RAINFAST: 4-hours
	HERBICIDE TANK MIXES: None registered

WEEDS SUPPRESSED

- Dandelion (>15 cm diameter)

MIXING INSTRUCTIONS

1. Fill the spray tank with 1/2 to 3/4 of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add the required amount of Eclipse XC A
4. Add the required amount of Eclipse XC B
5. Add any required adjuvant or surfactants
6. Complete filling the sprayer tank with water

Note: Do not mix undiluted herbicides in the chem-handler

CROP ROTATION

The year following application, fields can be seeded to:

- Barley
- Canola
- Flax
- Forage grasses
- Mustard
- Oats
- Peas*
- Rye (not under-seeded with legumes, clover or alfalfa)
- Summerfallow
- Wheat

GRAZING AND HARVEST

Allow 3 to 5 days after application before grazing treated areas.

* For pea rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140mm (5.5 inches) and annual rainfall must be greater than 175mm (6.9 inches)