



Enlist Duo™ provides the convenience of both 2,4-D choline and glyphosate in one formulation for control of grasses and broadleaf weeds including hard-to-control and resistant weeds.

WHY USE ENLIST DUO™ HERBICIDE?

- Enlist Duo™ herbicide with Colex-D™ technology is designed to land and stay on target with excellent performance on the toughest weeds
- Multiple modes of action deliver superior control: Groups 4 & 9
- Improved tank stability for a spray solution that stays mixed
- Spray solution does not adhere to sprayer components providing easy and fast sprayer cleanout

ENLIST E3® SOYBEANS (1.74 L/AC)

WEEDS CONTROLLED

- | | | | |
|----------------------------------|--------------------------------------|--|----------------------------------|
| • Annual sow thistle | • Corn spurry | • Knotweed (before 4-leaf) | • Roundleaf mallow ² |
| • Annual sunflower | • Cow cockle | • Kochia | • Russian pigweed |
| • Barnyard grass | • Daisy fleabane | • Lady's-thumb | • Russian thistle |
| • Biennial wormwood ² | • Dandelion | • Lamb's-quarters | • Shepherd's purse |
| • Blue lettuce ² | • Dog mustard | • Large crabgrass | • Smooth crabgrass |
| • Bluebur | • Eastern black flowering nightshade | • Leafy spurge ² | • Smooth pigweed |
| • Burdock (before 4-leaf) | • Fall panicum | • Mouse-eared chickweed ² | • Stinkweed |
| • Burdock ² | • False flax | • Mustards (except green tansy) | • Sweet clover |
| • Canada fleabane | • Field bindweed ³ | • Narrow-leaved hawk's-beard | • Tartary buckwheat |
| • Canada thistle ^{3,5} | • Field horsetail | • Nightflowering catchfly | • Velvetleaf |
| • Cleavers | • Field pepperglass | • Oak leaf goosefoot | • Vetch |
| • Cocklebur | • Flixweed | • Palmer amaranth ³ | • Volunteer barley |
| • Common chickweed | • Giant foxtail | • Pennsylvania smartweed | • Volunteer canola ¹ |
| • Common milkweed ^{3,4} | • Giant ragweed | • Perennial sow-thistle ^{3,5} | • Volunteer wheat |
| • Common plantain | • Goat's-beard | • Pineappleweed | • Waterhemp |
| • Common purslane | • Green foxtail | • Quackgrass | • Wild buckwheat |
| • Common ragweed | • Green smartweed | • Redroot pigweed | • Wild oats |
| • Common tansy | • Hairy galinsoga | | • Wild proso millet |
| | • Hedge bindweed | | • Wild radish |
| | • Hemp-nettle | | • Wild tomato |
| | • Hoary cress | | • Yellow nutsedge ^{3,6} |

¹ Including glyphosate-tolerant and Clearfield® canola varieties.

² Top growth control only.

³ Use two applications for best control. The 2nd application should be no later than the R2 stage (full flowering stage) of soybeans.

⁴ Milkweed: 15-60 cm in height and actively growing.

⁵ Canada thistle and perennial sowthistle: should be from the rosette stage to 50 cm in height and actively growing.

⁶ Yellow nutsedge: 5-15 cm in height and actively growing.

APPLICATION GUIDELINES

CROPS Enlist™ corn Enlist E3® soybeans Burndown before field corn, wheat (spring, winter, durum), barley, rye	PACKAGING	Case: 2 x 8.7 L Tote: 556.8 L
	RATES	Enlist™ field corn and Enlist E3® soybeans: 1.18 - 1.74 L/ac Burndown ahead of corn and cereal crops: 0.89 - 1.74 L/ac
	ACRES TREATED	1.74 L/ac rate: • 10 ac/case • 320 ac/tote
	WATER VOLUME	10-15 gal/ac
	TIMING	Crop stage: • Enlist corn: Up to V8 growth stage or 120 cm height • Enlist E3 soybeans: Up to R2 stage (full flowering) • Burndown before field corn, wheat (spring, winter, durum), barley, rye: Prior to planting or after planting (BUT BEFORE CROP EMERGENCE)
	RAINFAST	2 hours
	TANK MIX HERBICIDES	Consult the Enlist Product Use Guide available at EnlistCanada.ca .

Application information

On-Target Application Requirements

Droplet Size: Coarse to extremely coarse (ASAE S-572 Standard) to greatly reduce drift potential.

Boom Height: 60 cm or less

Wind: 3-16 km/hr. Do not spray during a temperature inversion. Do not spray in winds that exceed 25 km/h

Enlist corn, Enlist E3 soybeans: Make 1 to 2 applications with a minimum of 12 days between applications.

Crop Rotation

Any crop may be grown the year following an application of Enlist Duo

Pre-harvest Interval

Enlist E3 soybeans:

Do not harvest for forage or hay. Do not graze treated Enlist E3 soybeans

Enlist corn:

Do not permit lactating dairy animals to graze fields within 7 days after application.

Do not harvest forage or cut hay within 30 days after application. Withdraw meat animals from treated fields at least 3 days before slaughter.

AVAILABLE IN BULK