

Korrex™ II

HERBICIDE

Korrex™ II herbicide delivers superior pre-seed control of kochia, including Group 2 and 9 resistant biotypes, as well as 21 other tough broadleaf weeds.

WHY USE KORREX™ II?

- **Superior pre-seed control of kochia**, including Group 2 and Group 9 resistant biotypes.
- **Easy to tank-mix with any form of glyphosate.**
- **Broad-spectrum broadleaf weed control**, including dandelion, narrow-leaved hawk's beard, wild buckwheat, flixweed and stinkweed.
- Extended control of volunteer canola flushes.
- Effective solution for herbicide resistance management.
- Excellent solution for Canada thistle control with a fall application.

KORREX II + GLYPHOSATE AT 0.5 REL/AC

BROADLEAF WEEDS CONTROLLED

- Canada fleabane*²
- Chickweed*
- Cleavers*
- Common ragweed*
- Cow cockle*
- Flixweed*
- Hemp-nettle*
- Kochia*
- Lamb's-quarters*

- Narrow-leaved hawk's beard
- Redroot pigweed*
- Russian thistle
- Shepherd's purse*
- Smartweed*
- Stinkweed*
- Volunteer canola*¹
- Volunteer flax
- Wild buckwheat*
- Wild mustard*

GRASS WEEDS CONTROLLED

- Downy brome
- Giant foxtail
- Green foxtail
- Persian darnel
- Volunteer barley
- Volunteer wheat
- Wild oats

PERENNIAL WEEDS CONTROLLED

- Dandelion* (seedling, overwintered rosettes, mature plants up to 30 cm diameter)
- Perennial sow thistle³

WEEDS SUPRESSED

- Annual sow thistle
- Scentless chamomile

SPRING AND FALL SOILACTIVE™ EXTENDED CONTROL

- Canada fleabane
- Chickweed⁴
- Cleavers⁴
- Common ragweed
- Dandelion seedling
- Flixweed
- Hemp-nettle⁴
- Lamb's-quarters⁴
- Narrow-leaved hawk's beard⁴
- Redroot pigweed
- Scentless chamomile
- Shepherd's purse⁴
- Smartweed⁴
- Stinkweed⁴
- Volunteer Canola (except Clearfield*)
- Wild buckwheat
- Wild mustard

KORREX II + GLYPHOSATE AT 1 – 2.8 REL/AC

- Annual sow thistle
- Canada thistle
- Quackgrass (rosette stage)

*Weed controlled by multiple effective modes of action.

¹Including all herbicide-tolerant canola varieties.

²Less than 8 cm in height.

³Applications made at advanced stages will reduce effectiveness.

⁴Will not provide extended control of Group 2 resistant biotypes.

APPLICATION GUIDELINES

CROPS <i>Prior to cereal crops (spring or fall application):</i> Barley Durum wheat Oats Spring wheat Winter wheat	PACKAGING	Korrex™ II A herbicide: 1 x 0.45 kg jug Korrex™ II B herbicide: 1 x 7.76 L jug
	RATES	Spring Rate: <ul style="list-style-type: none"> • Korrex™ II A herbicide: 5.7 g/ac • Korrex™ II B herbicide: 97 mL/ac Fall Rate: <ul style="list-style-type: none"> • Korrex II A: 8.1 g/ac • Korrex II B: 139 mL/ac
	ACRES TREATED	Spring: 80 ac/case Fall: 56 ac/case
	WATER VOLUME	Ground 20–40 L/ac (5–10 US gal/ac)
	TIMING	Spring application: Prior to seeding (no later than 48 hours after seeding) Fall application: From after harvest to freeze up
	RAINFEST	30 minutes
	TANK-MIX HERBICIDES	Compatible with all forms of glyphosate



DOWNLOAD THE 2025 FIELD GUIDE APP FOR
PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.

CROP ROTATION

The year following a spring Korrex™ II herbicide application, fields can be seeded to:

- Barley
- Canola
- Chickpeas
- Corn
- Field beans
- Flax
- Lentils
- Mustard
- Peas
- Potatoes (except seed potatoes)
- Soybeans
- Summerfallow
- Sunflower
- Wheat

APPLICATION TIMING AND SEEDING

- Korrex™ II can be applied in the spring prior to seeding wheat, barley or oats and as an initial treatment in summerfallow.
- Korrex II applied in the fall (**after August 1**), can be seeded to winter wheat that fall. Spring wheat, durum, barley and oats can be seeded the following spring.
- Spring rate used in the fall will cover more acres without compromising control.

GRAZING AND HARVEST

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest treated crop within 60 days after application.

