

Korrex™ II

HERBICIDE

Korrex™ II herbicide delivers superior pre-seed control of kochia, including Group 2 and 9 resistant biotypes, as well as 21 other tough broadleaf weeds.

WHY USE KORREX™ II HERBICIDE?

- Kochia control. Superior pre-seed control of kochia, including Group 2 and Group 9 resistant biotypes.
- Flexible mixing options. Easy-mixing tank additive for any glyphosate.
- Broad-spectrum broadleaf weed control, including dandelion, narrow-leaved hawk's beard, wild buckwheat, flixweed and stinkweed.
- Extended control of volunteer canola flushes.
- Effective solution for herbicide resistance management.
- Excellent solution for Canada thistle control with a fall application.

KORREX™ II + GLYPHOSATE

Korrex II tank mixed with glyphosate such as VP480, at 0.5 REL/ac, provides effective pre-seed control of the following weeds:

BROADLEAF WEEDS CONTROLLED

- Annual sowthistle²
- Canada fleabane^{3,5}
- Chickweed*
- Cleavers*
- Common ragweed*
- Cow cockle*
- Flixweed*
- Hemp-nettle*
- Kochia*
- Lamb's-quarters*
- Narrow-leaved hawk's beard
- Redroot pigweed*
- Russian thistle
- Scentless chamomile²
- Shepherd's purse*
- Smartweed*
- Stinkweed*
- Volunteer canola*¹

- Volunteer flax
- Wild buckwheat*
- Wild mustard*

ANNUAL GRASSES CONTROLLED

- Downy brome
- Giant foxtail
- Green foxtail
- Persian darnel
- Volunteer barley

- Volunteer wheat
- Wild oats

PERENNIAL WEEDS CONTROLLED

- Dandelion*
(seedling, overwintered rosettes, mature plants up to 30 cm diameter)
- Perennial sow thistle⁴

SPRING AND FALL EXTENDED CONTROL:

- Canada fleabane
- Canola (except Clearfield⁵)
- Chickweed⁵
- Cleavers⁵
- Common ragweed
- Dandelion seedling
- Flixweed
- Hemp-nettle⁵
- Lamb's-quarters⁵
- Narrow-leaved hawk's beard⁵
- Redroot pigweed
- Scentless chamomile
- Shepherd's purse⁵
- Smartweed⁵
- Stinkweed⁵
- Wild buckwheat
- Wild mustard

Korrex II tank mixed with glyphosate at 1-2.8 REL/ac also provides effective control of the following weeds:

- Annual sow thistle
- Canada thistle (rosette stage)
- Quackgrass

*Weed controlled through multi-mode of action
1 Including all herbicide-tolerant canola varieties
2 Suppression only
3 Less than 8 cm in height

4 Applications made at advanced stages will reduce effectiveness

5 Will not provide extended control of Group 2 resistant biotypes

APPLICATION GUIDELINES

<p>CROPS</p> <p><i>Prior to cereal crops (spring or fall application):</i></p> <p>Barley</p> <p>Durum wheat</p> <p>Oats</p> <p>Spring wheat</p> <p>Winter wheat</p>	<p>PACKAGING</p> <ul style="list-style-type: none"> • Korrex II A: 1 x 0.45 kg jug • Korrex II B: 1 x 7.76 L jug
	<p>RATES</p> <p>Spring Rate:</p> <ul style="list-style-type: none"> • Korrex II A: 5.7 g/ac • Korrex II B: 97 ml/ac <p>Fall Rate:</p> <ul style="list-style-type: none"> • Korrex II A: 8.1 g/ac • Korrex II B: 139 ml/ac
	<p>ACRES TREATED</p> <p>Spring: 80 ac/case Fall: 56 ac/case</p>
	<p>WATER VOLUME</p> <p>Ground 20-40 L/ac (5-10 gal/ac)</p>
	<p>TIMING</p> <p>Spring application: Prior to seeding (no later than 48 hours after seeding) Fall application: From after harvest to freeze up</p>
	<p>RAINFAST</p> <p>30 minutes</p>
	<p>HERBICIDE TANK MIX HERBICIDES</p> <ul style="list-style-type: none"> • VP480 • Compatible with all forms of glyphosate

MIXING INSTRUCTIONS

1. Fill sprayer tank ½ full of water
2. Start sprayer tank agitation
3. Add the required amount of Korrex II A Herbicide, continue agitation
4. Add the required amount of Korrex II B Herbicide, continue agitation
5. Add the required amount of glyphosate, continue agitation
6. Fill the sprayer tank with sufficient water to spray 20-40 L/ac (5-10 US gal/ac)

Note: Do not mix undiluted herbicides in the chem-handler

CROP ROTATION

The year following a spring Korrex II application, fields can be seeded to:

- Barley
- Canola
- Chickpeas
- Corn
- Field beans
- Flax
- Lentils
- Mustard
- Peas
- Potatoes (except seed potatoes)
- Soybeans
- Summerfallow
- Sunflower
- Wheat

APPLICATION TIMING AND SEEDING

- Korrex II can be applied in the spring prior to seeding wheat, barley or oats and as an initial treatment in summerfallow.
- Korrex II applied **after August 1** can be seeded to winter wheat that fall or spring wheat, durum, barley and oats the following spring.

GRAZING AND HARVEST

- Livestock may be grazed on treated crops 7 days following application
- Do not harvest treated crop within 60 days after application.