

Liquid Achieve™ SC

Better value control of
grass weeds in barley
and wheat.

HERBICIDE

WHY USE LIQUID ACHIEVE™ SC HERBICIDE?

- Effective control of wild oats, persian darnel, barnyard grass and green/yellow foxtail in wheat and barley.
- Wide window of application.
- Trusted crop safety.
- Multiple pack sizes: 80 acre cases or 480 acre drums.

WEEDS CONTROLLED

- Wild oats
- Persian darnel
- Barnyard grass
- Green foxtail
- Volunteer oats
- Yellow foxtail

CARRIER® ADJUVANT REQUIRED AT 0.5% V/V:

Carrier adjuvant technology now available for use with Liquid Achieve™ SC herbicide.

	4,542 Litre Tank (1,200 US Gallon Tank)	3,785 Litre Tank (1,000 US Gallon Tank)	3,028 Litre Tank (800 US Gallon Tank)
0.5% v/v	22.71 L of Carrier required (0.005 x 4,542 L) 3 jugs	18.93 L of Carrier required (0.005 x 3,785 L) 2.5 jugs	15.14 L of Carrier required (0.005 x 3,028 L) 2 jugs

LIQUID ACHIEVE SC AND CARRIER RATE CHARTS:

	Liquid Achieve SC Rate 200 ml/ac 8 L jug or 96 L drum		Carrier Rate 0.5% v/v 8 L jug
	40 ac/jug	480 ac/drum	
Water volume	4,542 Litre Tank (1,200 US Gallon Tank)		
5 US gal/ac	240 acres	6 jugs	0.50 drum
10 US gal/ac	120 acres	3 jugs	0.25 drum
Water volume	3,785 Litre Tank (1,000 US Gallon Tank)		
5 US gal/ac	200 acres	5 jugs	0.42 drum
10 US gal/ac	100 acres	2.5 jugs	0.21 drum
Water volume	3,028 Litre Tank (800 US Gallon Tank)		
5 US gal/ac	160 acres	4 jugs	0.33 drum
10 US gal/ac	80 acres	2 jugs	0.17 drum

Intake™ Adjuvant can be used at 0.66% v/v if Carrier Adjuvant is not available. Liquid Achieve SC requires the addition of ammonia (Finish) with all EC broadleaf tank-mixes at 0.25% v/v + Carrier adjuvant at 0.5% v/v if bicarbonates > 400 ppm, add AMS to the tank first at 1% v/v.

APPLICATION GUIDELINES

<p>CROPS</p> <p>Barley Durum wheat Fall rye Spring rye Spring wheat Triticale Winter wheat</p> <p><i>Cereal crops underseeded to legume forages:</i></p> <p>Alfalfa Bird's foot trefoil Clover Sainfoin</p>	<p>PACKAGING</p> <p>Case: 2 x 8 L jugs Pallet: 5 x 96 L drums</p>
	<p>RATES</p> <p>200 ml/ac</p>
	<p>ACRES TREATED</p> <ul style="list-style-type: none"> • 40 ac/jug • 480 ac/drum
	<p>WATER VOLUME</p> <ul style="list-style-type: none"> • Ground: 20-40 L/ac (5-10 US gal/ac) • Aerial: 12-18 L/ac (3-5 gal/ac)
	<p>TIMING</p> <p>Crop stage: 2-leaf to just prior to flag leaf emergence Weed stage: Wild oats 1 to 6-leaf; other grasses 1 to 4-leaf</p>
	<p>RAINFAST</p> <p>1 hour</p>
	<p>TANK MIX HERBICIDES</p> <ul style="list-style-type: none"> • Attain™ XC • Buctril® M • Curtail® M • Infinity® • Infinity FX • Lontrel™ • MCPA Ester 600 • Mextrol® • OcTTain™ XL • Pardner™ • Pixxaro™ • Prestige™ XL • Prominex™ • Thumper® • Trophy® • 2,4-D Ester 700



DOWNLOAD THE 2022 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

Seed to all major crops the year following application.

GRAZING AND HARVEST

Allow 16 days after application before cutting hay or harvesting forage.

Mature crops may be harvested 60 days after application.

AVAILABLE IN BULK