

Lontrel™ XC

HERBICIDE

The most effective in-season thistle control available – right down to the roots.

WHY USE LONTREL™ XC HERBICIDE?

- Translocation throughout the plant for **effective control for both Canada thistle and sow thistle**.
- Up to a **73% thistle stand reduction** the year after application.
- Ability to choose a rate that matches the size of your thistle problem
- Improved, stronger, more convenient formulation.

BROADLEAF WEEDS CONTROLLED¹

- Common groundsel
- Ragweed
- Scentless chamomile
- Tufted vetch
- Vetch
- Volunteer alfalfa
- Volunteer soybean
- Wild buckwheat

PERENNIAL WEEDS CONTROLLED¹

- Alsike clover
- Canada thistle
- Perennial sow thistle (*top growth*)
- Red clover
- White clover

WEEDS SUPPRESSED

- Ox-eyed daisy
- Sheep sorrel

GRAZING AND HARVEST

- For canola, allow 3 to 5 days before grazing treated areas.
- For field corn, allow 40 days before grazing treated areas or feeding cattle with corn silage from treated areas
- For all other crops, no label restrictions on the grazing of crops or forages treated with Lontrel™ XC.

¹ The performance of Lontrel XC is rate dependent; control is achieved at the rate of 137 mL/ac (0.34 L/ha) or higher.

CLOPYRALID: 121 g/ac (13 ac/jug)
CLOPYRALID: 82 g/ac (20 ac/jug)

CLOPYRALID: 61 g/ac (26 ac/jug)
CLOPYRALID: 40 g/ac (40 ac/jug)

APPLICATION GUIDELINES

CROPS Cereals Corn Grasses Oilseeds Other (refer to product label)	PACKAGING	Case: 4 x 2.67 L	
	RATES & ACRES TREATED	<ul style="list-style-type: none"> • Acres: 13-40 ac/jug • Under cool or dry conditions, control may be severely reduced 	
	WATER VOLUME	<ul style="list-style-type: none"> • Ground 40-80 L/ac (10-21 US gal/ac) • Aerial not registered 	
	TIMING	Canola stage: 2 to 6-leaf Cereal stage: 3 to flag-leaf Corn stage: Emergence to 8-leaf (VE-V6) Weed stage: Canada thistle rosette to pre-bud	
	RAINFAST	4 hours	
	TANK-MIX HERBICIDES	<ul style="list-style-type: none"> • Ares™ SN herbicide • Amity™ WDG herbicide • Odyssey® • Poast® Ultra 	<ul style="list-style-type: none"> • Select® • Compatible with all forms of glyphosate



DOWNLOAD THE 2025 FIELD GUIDE APP FOR
PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.

CROP ROTATION

- Barley
- Canola
- Flax
- Forage grasses
- Mustard
- Oats
- Rye
- Summerfallow
- Wheat

Soybeans, field peas (crop rotation) – Fields previously treated with Lontrel™ XC herbicide up to 0.17 L/ha can be seeded after a minimum of 10 months to soybeans or field peas. Very dry soil conditions following application can result in a risk of injury to soybeans or field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive (less than 14 cm rainfall) in the year of application, delay seeding soybeans and field peas an additional 12 months (total 22 months following application).



Have you relied on Eclipse™ XC herbicide for excellent weed control in glyphosate tolerant canola/corn?

Consider tank-mixing Lontrel™ XC with your glyphosate for the same trusted control.

CLOPYRALID: 40 g/ac
GLYPHOSATE: 180 g/ac