

# Lontrel™ XC

## HERBICIDE

The most effective in-season thistle control available – right down to the roots. **Now registered in corn!**

### WHY USE LONTREL™ XC HERBICIDE?

- **Thistle control:** Translocation throughout the plant for effective control for both Canada thistle and sow thistle
- **Second year benefit:** up to a 73% thistle stand reduction the year after application
- **Flexibility:** choose a rate that matches the size of your thistle problem
- Improved, stronger, more convenient formulation
- **Now registered in Corn and Canola**

#### WEEDS CONTROLLED<sup>1</sup>

- Alsike clover
- Canada thistle
- Common groundsel
- Perennial sow thistle (*top growth*)
- Ragweed
- Red clover
- Scentless chamomile
- Tufted vetch

- Vetch
- Volunteer alfalfa
- White clover
- Wild buckwheat

#### WEEDS SUPPRESSED

- Ox-eyed daisy
- Sheep sorrel

#### GRAZING AND HARVEST

- For field corn, allow 40 days before grazing treated areas or feeding cattle with corn silage from treated areas
- For canola, allow 3 to 5 days before grazing treated areas.
- For all other crops, no label restrictions on the grazing of crops or forages treated with Lontrel™ XC herbicide.

<sup>1</sup> The performance of Lontrel XC is rate dependent; control is achieved at the rate of 37 ml/ac (.34 L/ha) or higher.

## APPLICATION GUIDELINES

<b>CROPS</b>  Cereals  <b>Corn - NEW</b>  Grasses  Oilseeds  Other <i>(refer to product label)</i>	<b>PACKAGING</b>	<b>Case:</b> 4 x 2.67 L
	<b>RATES &amp; ACRES TREATED</b>	<ul style="list-style-type: none"> <li>• 26-40 ac/jug</li> <li>• Under cool or dry conditions, control may be seriously reduced</li> </ul>
	<b>WATER VOLUME</b>	<ul style="list-style-type: none"> <li>• Ground 40-80 L/ac (10-21 US gal/ac)</li> <li>• Aerial not registered</li> </ul>
	<b>TIMING</b>	<b>Canola stage:</b> 2 to 6-leaf <b>Cereal stage:</b> 3 to flag leaf <b>Corn stage:</b> Emergence VE-V6 (8-leaf) <b>Weed stage:</b> Canada thistle rosette to pre-bud
	<b>RAINFAST</b>	4 hours
	<b>TANK MIX HERBICIDES</b>	Can be tank mixed with most commonly used grass weed herbicides, including: <ul style="list-style-type: none"> <li>• Ares™ SN</li> <li>• Amity™ WDG</li> <li>• Odyssey®</li> <li>• Poast® Ultra</li> <li>• Select®</li> <li>• VP480</li> <li>• Compatible with all forms of glyphosate</li> </ul>

### MIXING INSTRUCTIONS

1. Fill the spray tank with ½ to ¾ of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add any required water conditioners
4. Add any tank mix partners that are a dry formulation
5. Add the required amount of grassy weed tank mix partner
6. Add the required amount of Lontrel XC
7. Add any required adjuvant or surfactants
8. Complete filling the sprayer tank with water

*Note: Do not mix undiluted herbicides in the chem-handler*

### CROP ROTATION

The year following application, fields can be seeded to barley, canola, flax, forage grasses, mustard, oats, rye, wheat or can be summerfallowed.

*For field pea and soybean rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140 mm (5.5 inches) and annual rainfall must be greater than 175 mm (6.9 inches).*

*Note: For rotation cautions according to use rates, please see the product label.*