

OcTtain™ XL

Trusted, easy to use, high performance broadleaf weed control.

HERBICIDE

WHY USE OCTTAIN™ XL HERBICIDE?

- Performance and value. The standard for kochia (including Group 2 and 9 resistant), plus wild buckwheat and 25 other key broadleaf weeds.
- Ease of use. Available in 40-acre cases, 240-acre drums and 1,280-acre totes.
- Crop-safe and flexible. Excellent crop safety in barley, wheat and durum.
- Tank mix with Simplicity™ GoDRI™ (wheat), Liquid Achieve™ (barley) herbicides and other grass weed control products.

WEEDS CONTROLLED

- Annual sunflower
- Blue lettuce^{1,4}
- Bluebur
- Burdock
- Cleavers³
- Cocklebur
- Dandelion^{2,4}
- Docks⁴
- Dog mustard⁴
- Field bindweed^{1,4}
- Field horsetail¹
- Field peppergrass⁴
- Flixweed
- Goat's-beard
- Gumweed⁴
- Hairy galinsoga⁴
- Hedge bindweed⁴
- Hemp nettle
- Hoary cress¹
- Kochia³
- Lamb's-quarters
- Leafy spurge^{1,4}
- Mustards (*except green and grey tansy*)
- Oak-leaved goosefoot⁴
- Plantain
- Prickly lettuce
- Ragweed
- Redroot pigweed⁴
- Round-leaved mallow
- Russian thistle⁴
- Shepherd's purse
- Smartweed⁴
- Stinkweed
- Stork's-bill
- Sweet clover
- Tansy mustard⁴
- Tartary buckwheat⁴
- Vetch
- Volunteer canola⁵
- Volunteer flax
- Wild buckwheat
- Wild radish

WEEDS SUPPRESSED

- Annual sow thistle⁴
- Canada thistle^{1,4}
- Chickweed³
- Perennial sow thistle¹
- Redroot pigweed

¹ Top growth control only

² Spring rosettes

³ Including Group 2 resistant biotypes

⁴ Requires the addition of 86 ml/ac (2 oz/ac) of 2,4-D Ester 700

⁵ All herbicide-tolerant varieties

APPLICATION GUIDELINES

CROPS Barley Durum wheat Spring wheat Winter wheat	PACKAGING Case: 2 x 9 L jugs Pallet: 5 x 108 L drums Tote: 576 L
	RATES 450 ml/ac
	ACRES TREATED • 20 ac/jug • 240 ac/drum • 1280 ac/tote
	WATER VOLUME • Ground 20–40 L/ac (5–10 US gal/ac) • Aerial 12–20 L/ac (3–5 US gal/ac)
	TIMING Crop stage: 4-leaf to just prior to flag leaf emergence Weed stage: 1 to 6-leaf
	RAINFAST 1 hour
	TANK MIX HERBICIDES • Assert® • Everest® • Horizon® • Liquid Achieve™ SC • Puma® Super • Simplicity™ • Simplicity™ GoDRI™ • Traxos®



DOWNLOAD THE 2022 FIELD GUIDE APP FOR
PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

- Alfalfa
- Barley
- Canola
- Corn
- Dry beans
- Fababeans
- Flax
- Forage grasses
- Lentils
- Mustard
- Oats
- Peas
- Potatoes
- Rye
- Soybeans
- Sugar beets
- Sunflowers
- Wheat.

GRAZING AND HARVEST

- Allow 7 days after application before grazing lactating animals.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 30 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.

AVAILABLE IN BULK