

Paradigm™ PRE

Arylex™ active

HERBICIDE

Invest in your crop's yield with advanced, early weed control and sustainable herbicide resistance management. Shift your pre-seed Paradigm.



WHY USE PARADIGM™ PRE HERBICIDE WITH ARYLEX™ ACTIVE?

- Controls your toughest weeds, including Group 2 resistant cleavers and hemp-nettle.
- Controls small or large weeds in cool spring or fall conditions.
- Herbicide resistance management.
- Tank-mix with any form of glyphosate for your pre-seed burndown.
- Paradigm™ PRE's innovative GoDRI™ formulation comes in convenient packaging, making it easy to mix and handle.
- Provides extended control of volunteer canola flushes.

PARADIGM PRE AT 7.5 G/AC + GLYPHOSATE AT 0.5 REL/AC

SPRING RATE

BROADLEAF WEEDS CONTROLLED

- Canada fleabane*²
- Cleavers* (up to 9 whorls)
- Common chickweed* (up to 8 leaves)
- Common ragweed*²
- Flixweed*
- Hemp-nettle*
- Lamb's-quarters* (up to 8 leaves)
- Narrow-leaved hawk's beard
- Russian thistle
- Shepherd's purse*
- Smartweed* (up to 8 leaves)
- Stinkweed*
- Volunteer canola¹
- Volunteer flax (up to 15 cm)
- Wild buckwheat* (1-2 leaves)
- Wild mustard*

GRASS WEEDS CONTROLLED

- Barnyard grass
- Downy brome
- Giant foxtail
- Green foxtail
- Persian darnel
- Volunteer barley
- Volunteer wheat
- Wild oats

PERENNIAL WEEDS CONTROLLED

- Dandelion* (spring rosettes up to 15 cm in diameter)

WEEDS SUPPRESSED

- Kochia

SPRING AND FALL SOILACTIVE™ EXTENDED CONTROL:

- Canada fleabane
- Chickweed³
- Cleavers³
- Common ragweed
- Dandelion seedling
- Flixweed
- Hemp-nettle³
- Lamb's-quarters³
- Narrow-leaved hawk's beard³
- Redroot pigweed
- Scentless chamomile
- Shepherd's purse³
- Smartweed³
- Stinkweed³
- Volunteer canola (except Clearfield[®])
- Wild buckwheat
- Wild mustard

*Controlled by multiple effective modes of action.
1 Including all herbicide-tolerant canola varieties.
2 Less than 8 cm in height and including Group 2 resistant biotypes.

³ Will not provide extended control of Group 2 resistant biotypes.

APPLICATION GUIDELINES

CROPS Prior to cereal crops (spring or fall application): Barley Durum wheat Oats Spring wheat Winter wheat	PACKAGING	4 x 0.6 kg jugs
	RATES	Spring Rate: 7.5 g/ac Fall Rate: 10 g/ac
	ACRES TREATED	Spring Rate: 80 ac/jug (320 ac/case) Fall Rate: 60 ac/jug (240 ac/case)
	WATER VOLUME	Ground 20-40 L/ac (5-10 US gal/ac)
	TIMING	Spring Application: Prior to seeding (no later than 48 hours after seeding) Fall Application: From after harvest to freeze up
	RAINFAST	1 hour
	TANK-MIX HERBICIDES	Compatible with all forms of glyphosate



DOWNLOAD THE 2025 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.

CROP ROTATION

- 10 months:** all major crops, except lentils and chickpeas
- 22 months:** all major crops, including lentils and chickpeas

APPLICATION TIMING AND SEEDING

- Paradigm™ PRE herbicide can be applied in the spring prior to seeding wheat, barley or oats.
- Paradigm™ PRE applied **after August 1** can be seeded to winter wheat that fall or spring wheat, durum, oats and barley the following spring.
- Using spring rate in the fall will cover more acres with Paradigm PRE without compromising control, spray early and save.

GRAZING AND HARVEST

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.
- Do not cut the treated crop for hay or silage within 21 days after application.