

# PrePass™ XC

Nothing hits harder.  
Or lasts longer.

## HERBICIDE

### WHY USE PREPASS™ XC HERBICIDE?

- Confidence in performance in all conditions.
- SoilActive™ technology for extended control of key broadleaf weeds.
- **Time management.** Get the work done earlier – spring or fall.
- Serviced and supported through one company.
- **Convenience.** Glyphosate herbicide included for ease of use and compatibility.

#### BROADLEAF WEEDS CONTROLLED

- Annual sow thistle\*<sup>3</sup>
- Canada fleabane\*
- Chickweed\*
- Cleavers\*
- Cow cockle\*
- Crabgrass<sup>3</sup>
- Flixweed\*
- Hemp-nettle\*
- Kochia
- Lamb's-quarters
- Narrow-leaved hawk's beard\*

- Narrow-leaved vetch<sup>3</sup>
- Prickly lettuce\*<sup>3</sup>
- Ragweed\*
- Redroot pigweed\*
- Russian thistle
- Scentless chamomile\*
- Shepherd's purse\*
- Smartweed\*
- Stinkweed\*
- Volunteer canola
- Volunteer flax
- Wild buckwheat\*
- Wild mustard\*

#### GRASS WEEDS CONTROLLED

- Annual blue grass<sup>3</sup>
- Downy brome
- Foxtail barley<sup>2</sup>
- Giant foxtail
- Green foxtail
- Persian darnel
- Volunteer barley
- Volunteer wheat
- Wild oats

#### PERENNIAL WEEDS CONTROLLED

- Annual blue grass<sup>3</sup>
- Canada thistle<sup>3</sup>
- Dandelion\*<sup>1</sup>
- Quackgrass<sup>3</sup>
- Toadflax<sup>3</sup>

#### WEEDS SUPPRESSED

- Annual sow thistle
- Perennial sow thistle

#### SPRING AND FALL SOILACTIVE™ EXTENDED CONTROL

- Canada fleabane
- Volunteer canola (except Clearfield\*)

- Chickweed<sup>2</sup>
- Cleavers<sup>2</sup>
- Common ragweed
- Dandelion seedling
- Flixweed
- Hemp-nettle<sup>2</sup>

- Lamb's-quarters<sup>2</sup>
- Narrow-leaved hawk's beard<sup>2</sup>
- Redroot pigweed
- Scentless chamomile seedlings

- Shepherd's purse<sup>2</sup>
- Smartweed<sup>2</sup>
- Stinkweed<sup>2</sup>
- Wild buckwheat

\*Weed controlled through multi-effective mode of action  
1 Seedling, over-wintered rosettes, mature plants up to 30 cm in diameter

2 PrePass XC will not provide extended control of Group 2 resistant biotypes  
3 With the use of additional glyphosate; refer to the label for rates

## APPLICATION GUIDELINES

<b>CROPS</b> Prior to cereal crops (spring or fall application):  Barley  Durum wheat  Oats  Spring wheat  Winter wheat	<b>PACKAGING</b>	<b>Case:</b> • PrePass XC A: 1 x 1.6 L jug • PrePass XC B: 2 x 7.5 L jugs <b>Tote:</b> • PrePass XC A: 4 x 12 L jugs • PrePass XC B: 450 L tote
	<b>RATES</b>	<b>Spring &amp; Fall Rate:</b> • PrePass XC A: 40 ml/ac • PrePass XC B: 375 ml/ac
	<b>ACRES TREATED</b>	• 40 ac/case • 1,200 ac/tote
	<b>WATER VOLUME</b>	Ground 20-40 L/ac (5-10 US gal/ac)
	<b>TIMING</b>	<b>Spring application:</b> Prior to seeding (no later than 48 hours after seeding) <b>Chemfallow:</b> When weeds are actively growing, 1 to 4-leaf stage <b>Fall application:</b> From after harvest to freeze up
	<b>RAINFAST</b>	30 minutes
<b>TANK MIX HERBICIDES</b>	VP480 (PrePass XC B)	

### MIXING INSTRUCTIONS

1. Fill the spray tank with 1/2 to 3/4 of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add the required amount of PrePass XC A
4. Add the required amount of PrePass XC B
5. Complete filling the sprayer tank with water

Note: Do not mix undiluted herbicides in the chem-handler

### CROP ROTATION

The year following a spring PrePass™ XC herbicide application, fields can be seeded to:

- Alfalfa
- Barley
- Canola
- Chickpeas
- Corn
- Faba beans
- Field beans
- Flax
- Lentils
- Mustard
- Oats
- Peas
- Potatoes (except seed potatoes)
- Soybeans
- Summerfallow
- Sunflower
- Wheat

### APPLICATION TIMING AND SEEDING

- PrePass XC can be applied in the spring prior to seeding wheat, barley or oats
- PrePass XC applied to fields **after August 1** but prior to freeze up can be seeded in the fall to winter wheat and to wheat, barley, oats or summerfallow the following spring.

### GRAZING AND HARVEST

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.

AVAILABLE IN BULK