

# Prestige™ XL

Maximize potential.  
Every acre, clean.

## HERBICIDE

### WHY USE PRESTIGE™ XL HERBICIDE?

- Excellent control of thistles, cleavers, kochia and more.
- Your cleanest cereal crops, year after year.
- Multiple pack sizes: cases and drums.

*This Prestige XL formulation requires the addition of 0.25% v/v ammonia to be mixed with Simplicity™ GoDRI™. If Bicarbonates are >400 ppm, add AMS to the tank first at 2% v/v. Go to [PrestigeSimplicitySupport.Corteva.ca](https://www.corteva.com/pesticides) or call 1-800-667-3852 for further details.*

### 950 ml/ac (HIGH RATE)

#### WEEDS CONTROLLED

- Annual sow thistle
- Annual sunflower
- Burdock
- Canada thistle<sup>1</sup>
- Chickweed<sup>3,4</sup>
- Cleavers<sup>3</sup>
- Cocklebur
- Common groundsel
- Dandelion<sup>2</sup>
- Field horsetail<sup>5</sup>
- Flixweed
- Hemp-nettle
- Kochia<sup>3</sup>
- Lamb's-quarters
- Perennial sow thistle<sup>1</sup>
- Plantain<sup>5</sup>
- Prickly lettuce
- Ragweed
- Redroot pigweed
- Round-leaved mallow
- Russian pigweed
- Scentless chamomile
- Shepherd's purse

- Smartweed
- Stinkweed
- Stork's-bill
- Tartary buckwheat
- Vetch
- Volunteer canola<sup>6</sup>
- Volunteer flax
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild radish

### 710 ml/ac (LOW RATE)

#### WEEDS CONTROLLED

- Annual sunflower
- Burdock
- Canada thistle<sup>1</sup>  
*(low infestations)*
- Cleavers<sup>3</sup>
- Cocklebur
- Field horsetail<sup>5</sup>
- Flixweed
- Kochia<sup>3</sup>
- Lamb's-quarters
- Plantain<sup>5</sup>
- Prickly lettuce
- Ragweed
- Shepherd's purse
- Stinkweed
- Stork's-bill
- Vetch
- Volunteer flax
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild radish

#### WEEDS SUPPRESSED

- Volunteer canola<sup>6</sup>

<sup>1</sup> Season-long control, with some regrowth in the fall (top growth control)

<sup>2</sup> Spring rosettes only

<sup>3</sup> Including Group 2 resistant biotypes

<sup>4</sup> Controls or suppresses the population that is present at time of spraying but not subsequent flushes

<sup>5</sup> Top growth only

<sup>6</sup> All herbicide-tolerant canola varieties

FLUROXYPYR: 58 g/ac  
(Low Rate: 43 g/ac)

CLOPYRALID: 40 g/ac  
(Low Rate: 30 g/ac)

MCPA ESTER: 227 g/ac  
(Low Rate: 170 g/ac)

## APPLICATION GUIDELINES

<b>CROPS</b>  Barley  Canary seed  Durum wheat  Forage grasses <i>(see label for complete list.)</i>  Oats  Spring wheat  Winter wheat	<b>PACKAGING</b>	<b>Case:</b> 2 x 9.5 L jugs <b>Drum:</b> 113.6 L drum
	<b>RATES</b>	<b>Low rate:</b> 710 ml/ac <b>High rate:</b> 950 ml/ac
	<b>ACRES TREATED</b>	<b>Low rate:</b> • 13.5 ac/jug • 160 ac drum <b>High rate:</b> • 10 ac/jug • 120 ac/drum
	<b>WATER VOLUME</b>	• Ground 20–40 L/ac (5–10 US gal/ac) • Aerial 12–20 L/ac (3–5 US gal/ac)
	<b>TIMING</b>	<b>Crop stage:</b> 3-leaf to just prior to flag leaf emergence <b>Weed stage:</b> 1 to 6-leaf stage
	<b>RAINFAST</b>	4 hours
	<b>TANK MIX HERBICIDES</b>	• Axial® • Assert® • Everest® • Horizon • Liquid Achieve™ SC • Puma® • Simplicity™ • Simplicity GoDRI™



DOWNLOAD THE 2022 FIELD GUIDE APP FOR  
PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

### CROP ROTATION

- Barley
- Canola
- Flax
- Mustard
- Oats
- Peas<sup>1</sup>
- Rye
- Sugar beets
- Summerfallow
- Wheat

### GRAZING AND HARVEST

- Livestock may graze treated areas 7 days after application.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 7 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.

<sup>1</sup> For pea or soybean rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140 mm (5.5 inches) and annual rainfall must be greater than 175 mm (6.9 inches).

AVAILABLE IN BULK