

Prominex™

Arylex™ active

HERBICIDE

Prominex™ delivers exceptional control of annual AND perennial broadleaf weeds with the convenience of an all-in-one formulation.

WHY USE PROMINEX™ HERBICIDE IN WHEAT AND BARLEY?

- **Unmatched Weed Control.** Prominex provides control of both annual AND perennial broadleaf weeds.
- **Just GO.** Thanks to Arylex active, you can Just GO on small or large broadleaf weeds, in early or late crop staging, and even in cool or dry conditions.
- **Tank Mix Flexibility.** Choose whether to tank mix with a Group 1 grass herbicide in your wheat or barley and a Group 2 grass herbicide in your wheat.
- **Convenient Formulation.** All-in-one ME formulation allows for ultimate convenience and easy loading.
- **Expanded Group 4 Mode of Action Control.** Tank mix with MCPA Ester 600 or 2, 4-D Ester for expanded Group 4 broadleaf weed control.

Prominex + 235 ml/ac (5 oz./ac) MCPA Ester 600

GRASS WEEDS CONTROLLED

- Barnyard grass

ANNUAL BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Annual sow thistle
- Ball mustard
- Burdock
- Canada fleabane**
- Canada thistle
- Chickweed*
- Cleavers**
- Cocklebur
- Common ragweed**
- Cow cockle
- Dandelion

- False ragweed
- Flixweed*
- Giant ragweed**
- Hemp-nettle*
- Henbit
- Kochia**
- Lamb's-quarters
- Nightshade species (Eastern black, hairy and cutleaf)
- Plantain
- Prickly lettuce
- Redroot pigweed
- Round-leaved mallow
- Shepherd's purse*
- Stinkweed

- Stork's-bill
- Velvetleaf
- Vetch
- Volunteer alfalfa
- Volunteer canola (all herbicide tolerant varieties)
- Volunteer flax
- Wild buckwheat
- Wild mustard
- Wild radish

WEEDS SURPRESSED

- Canada thistle
- Field horsetail
- Perennial sow-thistle
- Smartweed

*Including Group 2 resistant biotypes

**Including Groups 2 & 9 resistant biotypes

For a complete weed list and specific weed staging, please refer to the Prominex RA label.

PROMINEX APPLICATION GUIDELINES

CROPS

Barley
Durum wheat
Spring wheat
Winter wheat

RATES AND ACRES TREATED

RATES:

– 414 ml/ac

ACRES TREATED:

– 40 ac/case (20 ac/jug)

PACKAGING:

– Case: 2 x 8.3 L jugs

WHEN TO APPLY

• 3-leaf to just prior to flag leaf emergence

RAINFAST


• 4 hours

HERBICIDE TANK MIXES

• MCPA Ester 600 • 2, 4-D Ester 700 • Simplicity™
• Simplicity™ GoDRI • Everest® • Axial®
• Axial + MCPA Ester 600
• Liquid Achieve™ SC • Trondus® • Horizon®
• Puma® Advance • Traxos®

2020 Research Authorizations

Prominex was showcased in 53 x 80 acre research authorization sites applied by farmers throughout Alberta, Saskatchewan and Manitoba.



Prominex + Simplicity GODRI (10 DAA)

Crop rotation

- Barley
- Canola
- Corn
- Flax
- Forage grasses
- Oats
- Mustard

- Peas¹
- Fall rye (*not underseeded with legumes, clover or alfalfa*)
- Soybeans
- Summerfallow
- Wheat

22 months:

- Lentils
- Chickpeas

For pea or soybean rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140 mm (5.5 inches) and annual rainfall must be greater than 175 mm (6.9 inches).