

Prospect™

Arylex™ active

Give your canola the best start possible with the true pre-seed weed control power of Prospect™ herbicide.

HERBICIDE

WHY USE PROSPECT™ HERBICIDE?

- Controls a wide range of broadleaf weeds, including cleavers (overwintered and Group 2 resistant biotypes), hemp-nettle, flixweed, narrow-leaved hawk's beard and many more.
- More consistent and complete control than glyphosate alone.
- Flexibility to spray at 5 gal/ac: low spray water volume without giving up weed control performance.
- Tank mixed with glyphosate, provides 3 modes of action for multi-mode of action control of key broadleaf weeds.

PROSPECT + GLYPHOSATE AT 0.5 REL/AC:

ANNUAL BROADLEAF WEEDS CONTROLLED

- American Dragonhead
- Canada fleabane*
- Chickweed
- Cleavers*
- Common ragweed*
- Cow Cockle
- Eastern black nightshade
- Flixweed
- Hemp-nettle*
- Henbit
- Kochia
- Lamb's-quarters
- Morning glory
- Narrow-leaved
- Redroot pigweed
- Round-leaved mallow
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Stork's bill
- Velvetleaf
- Volunteer canola (all herbicide tolerant varieties)
- Volunteer flax
- Waterhemp
- Wild buckwheat
- Wild mustard

hawk's beard

MULTI MODE OF ACTION

WEEDS:

- Barnyard grass
- Canada fleabane
- Chickweed
- Cleavers
- Eastern black nightshade
- Flixweed
- Hemp-nettle
- Lamb's-quarters
- Redroot pigweed
- Russian thistle
- Shepherd's-purse
- Smartweed
- Stinkweed
- Wild buckwheat
- Wild mustard

ANNUAL GRASS WEEDS CONTROLLED

- Barnyard grass
- Downy brome
- Giant foxtail
- Green foxtail
- Persian darnel
- Volunteer barley
- Volunteer wheat
- Wild oats

PERENNIAL WEEDS

- Dandelion
- Volunteer alfalfa

WEEDS SUPPRESSED

- Annual sow thistle

*Including group 2 resistant biotypes

APPLICATION GUIDELINES

CROPS <i>Prior to seeding:</i> Canola	PACKAGING	2 x 10.8 L jugs
	RATES	135 ml/ac
	ACRES TREATED	80 ac/jug (160 ac/case)
	WATER VOLUME	Ground 20-40 L/ac (5-10 gal/ac)
	TIMING	Prior to seeding canola
	RAINFAST	1 hour
	TANK MIX HERBICIDES	<ul style="list-style-type: none">• VP480• Compatible with all forms of glyphosate

MIXING INSTRUCTIONS

1. Fill sprayer tank 1/2 full of water
2. Start sprayer tank agitation
3. Add the required amount of Prospect herbicide, continue agitation
4. Add the required amount of glyphosate, continue agitation
5. Fill the sprayer tank with sufficient water to spray 20-40 L/ac (5-10 US gal/ac)

CROP ROTATION

10 months: All major crops, except lentils or chickpeas

22 months: Lentils, chickpeas or any other crops not listed

GRAZING AND HARVEST

Livestock may be grazed on treated crops 7 days following application.

Do not harvest the treated crop within 60 days after application.

APPLICATION TIMING

Prospect is registered for spring application only.