

# Prospect™

Arylex™ active

## HERBICIDE

Give your canola the best start possible.

### WHY USE PROSPECT™ HERBICIDE WITH ARYLEX™ ACTIVE?

- Controls a wide range of broadleaf weeds, including cleavers. (overwintered and Group 2 resistant biotypes), hemp-nettle, flixweed, wild buckwheat, and many more.
- More consistent and complete pre-seed control than glyphosate alone.
- Flexibility to spray at 5 gal/ac: low spray water volume without giving up weed control performance.\*
- Tank-mixed with glyphosate, provides 3 actives for multiple effective modes of action control of competitive weeds.

### PROSPECT™ + GLYPHOSATE AT 0.5 REL/AC:

#### BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Canada fleabane<sup>1</sup>
- Chickweed
- Cleavers<sup>1</sup>
- Common ragweed<sup>1</sup>
- Cow cockle
- Eastern black nightshade
- Flixweed
- Hemp-nettle<sup>1</sup>
- Henbit
- Kochia<sup>2</sup>
- Lamb's-quarters
- Morning glory

- Narrow-leaved hawk's beard<sup>3</sup>
- Redroot pigweed
- Round-leaved mallow
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Stork's-bill
- Velvetleaf
- Volunteer canola (all herbicide tolerant varieties)<sup>4</sup>
- Volunteer flax
- Waterhemp
- Wild buckwheat
- Wild mustard

#### WEEDS CONTROLLED THROUGH MULTIPLE EFFECTIVE MODES OF ACTION

- Barnyard grass
- Canada fleabane
- Chickweed
- Cleavers
- Eastern black nightshade

#### GRASS WEEDS CONTROLLED

- Barnyard grass
- Downy brome
- Giant foxtail
- Green foxtail
- Persian darnel
- Volunteer barley
- Volunteer wheat
- Wild oats

#### PERENNIAL WEEDS CONTROLLED

- Dandelion
- Volunteer alfalfa

#### WEEDS SUPPRESSED

- Annual sow thistle

\* Higher water volumes promote better coverage for heavy infestation or hard to control weeds.  
<sup>1</sup> Including Group 2 resistant biotypes.  
<sup>2</sup> Control of light to moderate infestation ≤150 plants/m<sup>2</sup>, ≤15cm in height, suppression of Group 9 resistant kochia biotypes.

<sup>3</sup> Up to 8cm in height; for larger stages use additional glyphosate top-up rates (see label).  
<sup>4</sup> Group 9 resistant varieties: under conditions of high density or larger stages control may be reduced.  
<sup>5</sup> Seeding/planting depth: minimum 4 cm (1.6 inches) or injury may occur.

### APPLICATION GUIDELINES

<b>CROPS</b> Prior to seeding:  Canola  Corn <sup>5</sup>	<b>PACKAGING</b>	2 x 10.8 L jugs
	<b>RATES</b>	135 mL/ac
	<b>ACRES TREATED</b>	80 ac/jug (160 ac/case)
	<b>WATER VOLUME</b>	Ground 20–40 L/ac (5–10 US gal/ac)
	<b>TIMING</b>	Prior to seeding
	<b>RAINFAST</b>	1 hour
	<b>TANK-MIX HERBICIDES</b>	Compatible with all forms of glyphosate



DOWNLOAD THE 2025 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.

### CROP ROTATION

**10 months:** All major crops, except lentils or chickpeas  
**22 months:** Lentils, chickpeas or any other crops not listed

### APPLICATION TIMING

Prospect can be applied in the spring, prior to seeding canola or corn.

### GRAZING AND HARVEST

Livestock may be grazed on treated crops 7 days following application.

### PRE-HARVEST INTERVAL

60 days

