

Rexade™

Arylex™ active

HERBICIDE

The complete wheat herbicide.

GoDRI
FORMULATION

WHY USE REXADE™ HERBICIDE WITH ARYLEX™ ACTIVE?

- **Exceptional** grass and broadleaf weed control.
- **Complete, convenient**, all-in-one box solution.
- GoDRI™ Rapid Dispersion Technology for easy storage, transport, and mixing.

BROADLEAF
WEEDS
CONTROLLED

- American dragonhead
- Annual sunflower
- Bluebur
- Burdock
- Canada fleabane⁵
- Chickweed²
- Cleavers³
- Cocklebur
- Common ragweed⁴
- Corn spurry
- Cow cockle²
- Flixweed²
- Hemp-nettle^{2,3}
- Henbit⁴
- Lamb's-quarters^{2,3}
- Mustard² (except dog and green tansy)

- Plantain
- Prickly lettuce
- Redroot pigweed^{2,3}
- Round-leaved mallow²
- Russian thistle¹
- Shepherd's purse²
- Smartweed
- Stinkweed²
- Stork's-bill
- Sweet clover
- Velvetleaf
- Volunteer alfalfa
- Volunteer canola
- Volunteer flax
- White cockle
- Wild buckwheat²
- Wild radish¹

GRASS WEEDS
CONTROLLED

- Barnyard grass
- Japanese brome
- Wild oats
- Yellow foxtail

WEEDS
SUPPRESSED

- Annual sow thistle
- Canada thistle
- Dandelion
- Downy brome
- Green foxtail
- Kochia^{1,3}
- Night-flowering catchfly

1 2 to 4-leaf staging.
2 Controlled through multiple effective modes of action.

3 Including Group 2 resistant biotypes.
4 Up to bud stage and 15 cm in height.
5 Including Group 2 and Group 9 resistant biotypes.

APPLICATION GUIDELINES

CROPS Durum wheat Spring wheat Winter wheat	PACKAGING	• Rexade™ A: 1 x 1.62 kg jug • Rexade™ B: 1 x 8.58 L jug
	RATES	• Rexade™ A: 40.5 g/ac • Rexade™ B: 215 mL/ac
	ACRES TREATED	40 ac/case
	TIMING	Crop stage: 2 leaf (fully expanded) to flag leaf
	RAINFAST	2 hours



DOWNLOAD THE 2025 FIELD GUIDE APP FOR
PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.

CROP ROTATION

- Barley
 - Brown mustard
 - Canola
 - Peas
- Flax
 - Wheat
 - Oats
 - Soybeans
- Sunflowers
 - Potatoes (except seed potatoes)
 - Yellow mustard

GRAZING AND HARVEST

- Allow 7 days after application before grazing lactating animals.
- Livestock may graze treated areas 3 days after application.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 30 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.

