

Rexade™

Arylex™ active

HERBICIDE

With Arylex™ active Just GO benefits, controls all your standard broadleaf weeds in wheat.

**WHY USE REXADE™ HERBICIDE IN WHEAT?**

- No compromise. Unmatched cross-spectrum grass and broadleaf weed control in one product.
- New class of Group 4 broadleaf herbicide combined with a premium Group 2 grass herbicide.
- Convenience. Complete, all-in-one box solution.
- GoDRI™ Rapid Dispersion Technology for easy storage, transport and mixing.
- Multiple Mode of Action control of key weeds for effective resistance management.

GRASS WEEDS CONTROLLED

- Wild oats
- Barnyard grass
- Japanese brome
- Yellow foxtail

BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Annual sunflower
- Bluebur
- Burdock
- Canada fleabane
- Chickweed**
- Cleavers¹
- Cocklebur
- Common ragweed²
- Corn spurry
- Cow cockle**
- Flixweed**
- Hemp-nettle**¹
- Henbit²
- Kochia^{1*}
- Lamb's-quarters**¹
- Mustard** (*except dog and green tansy*)
- Plantain
- Prickly lettuce
- Redroot pigweed**¹
- Round-leaved mallow**
- Russian thistle*
- Shepherd's purse**
- Smartweed
- Stinkweed**
- Stork's-bill
- Sweet clover
- Velvetleaf
- Volunteer alfalfa
- Volunteer canola
- Volunteer flax
- White cockle
- Wild buckwheat**
- Wild radish*

WEEDS SUPPRESSED

- Annual sow thistle
- Canada thistle
- Dandelion
- Downy brome
- Green foxtail
- Night-flowering catchfly

*2 to 4-leaf staging

**Weed controlled through multi-mode of action

¹ Including Group 2 resistant biotypes² Up to bud stage and 15 cm in height³ Including Group 2 and Group 9 resistant

APPLICATION GUIDELINES

CROPS Durum wheat Spring wheat Winter wheat	PACKAGING <ul style="list-style-type: none"> • Rexade A: 1 x 1.62 kg jug • Rexade B: 1 x 8.58 L jug
	RATES <ul style="list-style-type: none"> • Rexade A: 40.5 g/ac • Rexade B: 215 ml/ac
	ACRES TREATED <p>40 ac/case</p>
	TIMING <p>Crop stage: 2-leaf to just prior to flag leaf emergence</p>
	RAINFAST <p>2 hours</p>

MIXING INSTRUCTIONS

1. Fill the spray tank with ½ to ¾ of the required amount of water
2. Continue agitation throughout the mixing and spraying procedure
3. Add the required amount of Rexade A
4. Add the required amount of Rexade B
5. Complete filling the sprayer tank with water

Note: Do not mix undiluted herbicides in the chem-handler

No tank mix purchase necessary, everything included in the box

CROP ROTATION

All major crops the following spring, except lentils or chickpeas.

GRAZING AND HARVEST

- Allow 7 days after application before grazing lactating animals.
- Livestock may graze treated areas 3 days after application.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 30 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.