

# Simplicity™ GoDRI™

## HERBICIDE

Superior performance, including elite Group 2 wild oat and bonus broadleaf weed control with no re-cropping restrictions in a convenient, easy-to-use GoDRI formulation.

**GoDRI**  
FORMULATION

## WHY USE SIMPLICITY™ GODRI™ HERBICIDE?

- Superior performance. Elite grass and broadleaf weed control.
- Wide window of application. This is the only Group 2 wild oat product that can be applied up to just prior to flag leaf emergence.
- Resistance management. Controls Group 1 resistant wild oats.
- Broadleaf herbicide tank mix partner flexibility.
- Rotational freedom to all major crops the following year.
- Convenient GoDRI™ herbicide formulation for fast, easy mixing and handling.

### GRASS WEEDS CONTROLLED

- Wild oats
- Barnyard grass
- Downy brome  
(fall application)
- Japanese brome
- Yellow foxtail

### BROADLEAF WEEDS CONTROLLED

- Chickweed
- Cleavers\*
- Corn spurry
- Cow cockle
- Flixweed
- Hemp-nettle
- Redroot pigweed
- Round-leaved mallow

- Shepherd's purse
- Smartweed\*
- Stinkweed
- Volunteer canola  
(excluding Clearfield®)

### WEEDS SUPPRESSED

- Canada thistle
- Dandelion
- Downy brome  
(spring application)
- Green foxtail<sup>1</sup>
- Persian darnel
- Russian thistle
- White cockle
- Wild buckwheat

**When tank mixing Simplicity GoDRI with a broadleaf herbicide that does not require a surfactant, Bindem Utility Modifier must be used. Bindem is sold separately.**

**When spraying Simplicity™ GoDRI™ alone, the addition of a non-ionic surfactant is ALWAYS required. The following non-ionic surfactants can be used:**

Agral 90 at 0.25% v/v, Sentry at 0.25% v/v, Ag-Surf® Original at 0.25% v/v

\*Not including group 2 resistant

<sup>1</sup> Corteva Agriscience research trials indicate that application to small stage, actively growing plants provides an increased level of control

APPLICATION GUIDELINES

|  |                                    |  |
|--|------------------------------------|--|
| <p><b>CROPS</b></p> <p>Durum wheat</p> <p>Spring wheat</p> <p>Winter wheat</p> | <p><b>PACKAGING</b></p>            | 4 x 2.24 kg jugs   |
|  | <p><b>RATES</b></p>                | 28 g/ac  |
|  | <p><b>ACRES TREATED</b></p>        | 80 ac/jug (320 ac/case)  |
|  | <p><b>WATER VOLUME</b></p>         | <ul style="list-style-type: none"> <li>• Ground 12-40 L/ac (3-10 US gal/ac)</li> <li>• Aerial minimum 12 L/ac (3 US gal/ac)</li> </ul>   |
|  | <p><b>TIMING</b></p>               | <p><b>Crop stage:</b> 2-leaf to just prior to flag leaf emergence</p> <p><b>Weed stage:</b></p> <ul style="list-style-type: none"> <li>• 1 to 4-leaf, two tillers stage of wild oats</li> <li>• 1 to 5-leaf stage of broadleaf weeds</li> </ul>  |
|  | <p><b>RAINFAST</b></p>             | 2 hours  |
|  | <p><b>TANK MIX HERBICIDES*</b></p> | <ul style="list-style-type: none"> <li>• Attain™ XC</li> <li>• Buctril® M</li> <li>• Cirpreme™ XC</li> <li>• Curtail® M</li> <li>• Exhilarate™</li> <li>• MCPA Ester 600</li> <li>• OctTain™ XL</li> <li>• Prestige™ XL</li> <li>• Prominex™</li> <li>• Pixxaro™</li> <li>• Refine Extra®</li> <li>• Stellar™ XL</li> <li>• Thumper®</li> <li>• Tilt® fungicide</li> </ul> |



DOWNLOAD THE 2022 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

**CROP ROTATION**

- Alfalfa
- Barley
- Brown mustard
- Canola
- Canola quality brassica juncea
- Chickpea
- Dry bean (*of the species phaseolus vulgaris*)
- Field corn
- Field peas
- Flax
- Lentils
- Oats
- Potatoes
- Soybean
- Spring wheat
- Sunflower
- Yellow mustard

**GRAZING AND HARVEST**

- Livestock may graze treated areas 7 days after application.
- Mature crops may be harvested 60 days after application.

^ Unless otherwise noted.  
Always read and follow label directions.