

# Simplicity™ GoDRI™

## HERBICIDE

Simplicity™ GoDRI™ herbicide Wild Oat Rate. Simply better value.

**GODRI**  
FORMULATION

### WHY USE SIMPLICITY™ GODRI™ WILD OAT RATE?


- **Cost effective control of wild oats** and Japanese brome.
- **Wide window of application** from 2-leaf to flag-leaf.
- **Rotational freedom** - ability to seed all major crops the year following application.
- Convenient GoDRI™ herbicide formulation for fast, easy mixing and handling.

For early season applications in low to moderate wild oat populations that are typical in the brown soil zones of Southern Alberta and Southern Saskatchewan.

### WILD OAT RATE GRASS WEEDS CONTROLLED

- Wild oats
- Japanese brome<sup>1</sup>
- Barnyard grass<sup>1</sup>

### WILD OAT PERFORMANCE



**WHEN TANK-MIXING SIMPLICITY GODRI WITH A BROADLEAF HERBICIDE THAT DOES NOT REQUIRE A SURFACTANT, BINDEM UTILITY MODIFIER MUST BE USED. BINDEM IS SOLD SEPARATELY.**

When spraying Simplicity GoDRI alone, the addition of a non-ionic surfactant is ALWAYS required. The following non-ionic surfactants can be used: Agral 90 at 0.25% v/v, Sentry at 0.25% v/v, Ag-Surf® Original at 0.25% v/v

<sup>1</sup>Based on Corteva Agriscience research trials.

### APPLICATION GUIDELINES

<b>CROPS</b> Durum wheat Spring wheat Winter wheat Fall Rye Spring Rye Triticale	<b>PACKAGING</b>	4 x 2.24 kg jugs
	<b>RATES</b>	21 g/ac
	<b>ACRES TREATED</b>	106 ac/jug (424 ac/case)
	<b>WATER VOLUME</b>	<ul style="list-style-type: none"> <li>• Ground 20-40 L/ac (5-10 US gal/ac)</li> <li>• Aerial minimum 12 L/ac (3 US gal/ac)</li> </ul>
	<b>TIMING</b>	<b>Crop stage:</b> 2 leaf (fully expanded) to flag-leaf <b>Weed stage:</b> 1 to 4-leaf
	<b>RAINFAST</b>	2 hours
	<b>TANK-MIX HERBICIDES<sup>2</sup></b>	<ul style="list-style-type: none"> <li>• Attain™ XC herbicide</li> <li>• Buctril® M</li> <li>• OcTTain™ XL herbicide</li> <li>• Prestige™ XL herbicide</li> <li>• Prominex™ herbicide</li> <li>• Stellar™ XL herbicide</li> <li>• Thumper®</li> <li>• Tilt® fungicide</li> </ul>



**DOWNLOAD THE 2025 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.**

### CROP ROTATION

- Alfalfa
- Barley
- Brown mustard
- Canola
- Canola-quality Brassica juncea
- Chickpea
- Dry bean (*Phaseolus vulgaris* species)
- Field corn
- Field peas
- Flax
- Lentils
- Oats
- Potatoes
- Soybean
- Spring wheat
- Sunflower
- Yellow mustard

### GRAZING AND HARVEST

- Livestock may graze treated areas 7 days after application.
- Mature crops may be harvested 50 days after application.



<sup>2</sup> Unless otherwise stated. Always read and follow label directions.