

Tandem™

HERBICIDE

Control problem grass and broadleaf weeds in wheat – with the tougher, easier, total-acre solution.

WHY USE TANDEM™ HERBICIDE?

- Tougher. Control of wild oats, Japanese brome, chickweed, cleavers, hemp-nettle, kochia, wild buckwheat and many more.
- Easy to use. Control all problem weeds using one solution.
- Flexible. Wide window of application, excellent crop safety, flexible rate and rotational freedom.
- Multi-Mode of Action. Two unique modes of action provide overlapping activity on hard-to-kill weeds for superior resistance management.

TANDEM ALONE

WEEDS CONTROLLED

GRASS

- Barnyard grass
- Japanese brome
- Wild oats¹
- Yellow foxtail

BROADLEAF

- Chickweed^{2*}
- Cleavers^{2*}
- Corn spurry
- Cow cockle

- Flixweed
- Hemp-nettle
- Kochia²
- Redroot pigweed
- Round-leaved mallow*
- Shepherd's purse
- Smartweed
- Stinkweed
- Volunteer canola**
- Volunteer flax

WEEDS SUPPRESSED

GRASS

- Downy brome⁴
- Green foxtail³

BROADLEAF

- Canada thistle
- Dandelion⁴
- Russian thistle
- Stork's-bill
- Wild buckwheat

TANDEM + 241 ml/ac OF 2,4-D ESTER 700 (6 oz/ac)

ALL TANDEM ALONE

WEEDS PLUS:

- Bluebur
- Burdock
- Cocklebur

- Flixweed*
- Goat's-beard
- Lamb's-quarters*
- Plantain
- Prickly lettuce
- Ragweed
- Redroot pigweed*
- Smartweed*
- Stinkweed*
- Sweet clover

- Volunteer sunflower
- Wild buckwheat
- Wild mustard*
- Wild radish

TANDEM + 234 ml/ac OF MCPA ESTER 600 (5 oz/ac)

ALL TANDEM ALONE

WEEDS PLUS:

- Burdock
- Cocklebur
- Lamb's-quarters*

- Prickly lettuce
- Ragweed
- Shepherd's purse*
- Smartweed*
- Stinkweed*
- Vetch
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild radish

¹Weed controlled through multi-mode of action

²Clearfield volunteer canola is controlled with the addition of MCPA or 2,4-D

³Including Group 1 resistant biotypes

⁴Including Group 2 resistant biotypes

³Corteva Agriscience research trials indicate that application to small stage, actively growing plants provides an increased level of control

⁴Spring rosettes

⁵Top growth only

APPLICATION GUIDELINES

| | | |
|--------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>CROPS</p> <p>Durum wheat</p> <p>Spring wheat</p> <p>Winter wheat</p> | <p>PACKAGING</p> | <p>Case:</p> <ul style="list-style-type: none"> • Tandem A: 1 x 8 L jug • Tandem B: 1 x 4.84 L jug • Water conditioner: 1 x 1.5 L jug |
| | <p>RATES</p> | <ul style="list-style-type: none"> • Tandem A: 200 ml/ac • Tandem B: 121 ml/ac • Water conditioner: 37.5 ml/ac |
| | <p>ACRES TREATED</p> | <p>40 ac/case</p> |
| | <p>WATER VOLUME</p> | <p>Ground: 20-40 L/ac (5-10 US gal/ac)</p> <p>Aerial: Min. 12 L/ac (3 US gal/ac)</p> |
| | <p>TIMING</p> | <p>Crop stage: 3-leaf to just prior to flag leaf emergence</p> <p>Weed stage: 1 to 4-leaf, two tillers stage for wild oats and 1 to 5-leaf stage for broadleaf weeds</p> |
| | <p>RAINFAST</p> | <p>2 hours</p> |
| <p>TANK MIX HERBICIDES</p> | <ul style="list-style-type: none"> • Curtail® M herbicide • MCPA Ester 600 • 2,4-D Ester 700 | |



DOWNLOAD THE 2022 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

- Barley
- Canola
- Corn
- Dry beans
- Flax
- Lentils
- Mustard
- Oats
- Peas
- Potatoes
- Soybeans
- Sunflower
- Wheat

GRAZING AND HARVEST

- Livestock may graze treated areas 7 days after application.
- Allow 7 days after application before grazing lactating animals.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 30 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.

Additional broadleaf weeds controlled when tank mixed with Curtail M herbicide (10 ac/jug)

- Annual sow thistle
- Burdock
- Canada thistle
- Cocklebur
- Common groundsel
- Dandelion⁴
- Field horsetail⁵
- Lamb's-quarters
- Perennial sow thistle
- Plantain⁶
- Prickly lettuce
- Ragweed
- Russian pigweed
- Scentless chamomile
- Sunflower (annual and volunteer)
- Tartary buckwheat
- Vetch
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild radish